

RC

268.5

U55

no. 142 National Cancer Institute

1979 **CARCINOGENESIS**

Technical Report Series

No. 142

1979

**BIOASSAY OF  
p-CRESIDINE  
FOR POSSIBLE CARCINOGENICITY**

**CAS No. 120-71-8**

**NCI-CG-TR-142**

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE  
Public Health Service  
National Institutes of Health



Library  
Product Analysis of Health  
Literacy Research 2014

BIOASSAY OF  
p-CRESIDINE  
FOR POSSIBLE CARCINOGENICITY

*United States Food and Drug Administration  
cc  
Carcinogenesis Technical Report*

Carcinogenesis Testing Program  
Division of Cancer Cause and Prevention  
National Cancer Institute  
National Institutes of Health  
Bethesda, Maryland 20014

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE  
Public Health Service  
National Institutes of Health

DHEW Publication No. (NIH) 79-1397

270  
2687  
125  
60,142  
1719

REPORT ON THE BIOASSAY OF p-CRESIDINE  
FOR POSSIBLE CARCINOGENICITY

CARCINOGENESIS TESTING PROGRAM  
DIVISION OF CANCER CAUSE AND PREVENTION  
NATIONAL CANCER INSTITUTE, NATIONAL INSTITUTES OF HEALTH

FOREWORD: This report presents the results of the bioassay of p-cresidine conducted for the Carcinogenesis Testing Program, Division of Cancer Cause and Prevention, National Cancer Institute (NCI), National Institutes of Health, Bethesda, Maryland. This is one of a series of experiments designed to determine whether selected chemicals have the capacity to produce cancer in animals. Negative results, in which the test animals do not have a significantly greater incidence of cancer than control animals, do not necessarily mean the test chemical is not a carcinogen because the experiments are conducted under a limited set of circumstances. Positive results demonstrate that the test chemical is carcinogenic for animals under the conditions of the test and indicate a potential risk to man. The actual determination of the risk to man from animal carcinogens requires a wider analysis.

CONTRIBUTORS: This bioassay of p-cresidine was conducted by Mason Research Institute, Worcester, Massachusetts, initially under direct contract to the NCI and currently under a subcontract to Tracor Jitco, Inc., prime contractor for the NCI Carcinogenesis Testing Program.

The experimental design was determined by the NCI Project Officers, Dr. J. H. Weisburger (1,2) and Dr. E. K. Weisburger (1). The principal investigators for the contract were Dr. E. Smith (3) and Dr. A. Handler (3). Animal treatment and observation were supervised by Mr. G. Wade (3) and Ms. E. Zepp (3). Chemical analysis was performed by Midwest Research Institute (4) and the analytical results were reviewed by Dr. N. Zimmerman (5).

Histopathologic examinations were performed by Dr. D. W. Hayden (3), Dr. A. S. Krishna Murthy (3) and Dr. A. Russfield (3) at the Mason Research Institute, and the diagnoses included in this report represent the interpretation of these pathologists. Histopathology findings and reports were reviewed by Dr. R. L. Schueler (6).

Compilation of individual animal survival, pathology, and summary tables was performed by EG&G Mason Research Institute (7); the statistical analysis was performed by Mr. W. W. Belew (5,8) and Mr. R. M. Helfand (5), using methods selected for the Carcinogenesis Testing Program by Dr. J. J. Gart (9).

This report was prepared at METREK, a Division of The MITRE Corporation (5) under the direction of the NCI. Those responsible for this report at METREK are the project coordinator, Dr. L. W. Thomas (5), task leader Dr. M. R. Kornreich (5,10), senior biologist Ms. P. Walker (5), biochemist, Mr. S. C. Drill (5), and technical editor Ms. P. A. Miller (5). The final report was reviewed by members of the participating organizations.

The following other scientists at the National Cancer Institute were responsible for evaluating the bioassay experiment, interpreting the results, and reporting the findings: Dr. K. C. Chu (1), Dr. C. Cueto, Jr. (1), Dr. J. F. Douglas (1), Dr. D. G. Goodman (1,10), Dr. R. A. Griesemer (1), Dr. M. H. Levitt (1), Dr. H. A. Milman (1), Dr. T. W. Orme (1), Dr. R. A. Squire (1,11), Dr. S. F. Stinson (1), Dr. J. M. Ward (1), and Dr. C. E. Whitmire (1).

- 
1. Carcinogenesis Testing Program, Division of Cancer Cause and Prevention, National Cancer Institute, National Institutes of Health, Bethesda, Maryland.
  2. Now with the Naylor Dana Institute for Disease Prevention, American Health Foundation, Hammon House Road, Valhalla, New York.
  3. Mason Research Institute, 57 Union Street, Worcester, Massachusetts.
  4. Midwest Research Institute, 425 Volker Boulevard, Kansas City, Missouri.
  5. The MITRE Corporation, METREK Division, 1820 Dolley Madison Boulevard, McLean, Virginia.
  6. Tracor Jitco, Inc., 1776 East Jefferson Street, Rockville, Maryland.
  7. EG&G Mason Research Institute, 1530 East Jefferson Street, Rockville, Maryland.
  8. Now with the Solar Energy Research Institute, Cole Boulevard, Golden, Colorado.
  9. Mathematical Statistics and Applied Mathematics Section, Biometry Branch, Field Studies and Statistics Program, Division of Cancer Cause and Prevention, National Cancer Institute, National Institutes of Health, Bethesda, Maryland.



10. Now with Clement Associates, Inc., 1010 Wisconsin Avenue, N.W., Washington, D.C.
11. Now with the Division of Comparative Medicine, Johns Hopkins University, School of Medicine, Traylor Building, Baltimore, Maryland.

THE UNIVERSITY OF CHICAGO

PHYSICS DEPARTMENT

PHYSICS 350

LECTURE 1

1.1

1.2

1.3

1.4

1.5

1.6



## SUMMARY

A bioassay of p-cresidine for possible carcinogenicity was conducted using Fischer 344 rats and B6C3F1 mice. p-Cresidine was administered in the feed, at either of two concentrations, to groups of 50 male and 50 female animals of each species. The dietary concentrations used in the chronic bioassay for low and high dose rats were 0.5 and 1.0 percent, respectively. The time-weighted average concentrations fed to low dose male, low dose female, high dose male and high dose female mice were 0.22, 0.22, 0.46, and 0.44 percent, respectively.

All dosed animals, except for high dose male mice, were administered p-cresidine in the diet for 104 weeks and observed for an additional period of up to 2 weeks. All high dose male mice were dead by the end of week 92. For each species, 50 animals of each sex were placed on test as controls and fed only the basal laboratory diet.

Mortality rates were dose-related for both sexes of both species. That incidences of certain tumors were higher in low dose than in high dose groups was probably due to accelerated mortality in the high dose groups.

In dosed rats of both sexes, statistically significant incidences of bladder carcinomas (combined incidences of papillary carcinomas, squamous-cell carcinomas, transitional-cell papillomas, transitional-cell carcinomas, and undifferentiated carcinomas) and olfactory neuroblastomas were observed. The combined incidence of neoplastic nodules of the liver, hepatocellular carcinomas, or mixed hepato/cholangio carcinomas was also significant in low dose male rats.

In both male and female dosed mice, the incidence of bladder carcinomas (combined incidence of carcinomas NOS, squamous-cell carcinomas, and transitional-cell carcinomas) was significant. The incidence of hepatocellular carcinomas was significant in dosed female mice.

Under the conditions of this bioassay, p-cresidine was carcinogenic to Fischer 344 rats, causing increased incidences of carcinomas and of papillomas of the urinary bladder in both sexes, increased incidences of olfactory neuroblastomas in both sexes, and of liver tumors in males. p-Cresidine was also carcinogenic in B6C3F1 mice, causing carcinomas of the urinary bladders in both sexes and hepatocellular carcinomas in females.

The first part of the paper is devoted to a discussion of the general principles of the theory of the structure of the atom. It is shown that the structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum. The structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum.

The second part of the paper is devoted to a discussion of the general principles of the theory of the structure of the atom. It is shown that the structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum. The structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum.

The third part of the paper is devoted to a discussion of the general principles of the theory of the structure of the atom. It is shown that the structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum. The structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum.

The fourth part of the paper is devoted to a discussion of the general principles of the theory of the structure of the atom. It is shown that the structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum. The structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum.

The fifth part of the paper is devoted to a discussion of the general principles of the theory of the structure of the atom. It is shown that the structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum. The structure of the atom is determined by the laws of quantum mechanics, which are based on the principle of the conservation of energy and the principle of the conservation of momentum.

## TABLE OF CONTENTS

	<u>Page</u>
I. INTRODUCTION	1
II. MATERIALS AND METHODS	4
A. Chemicals	4
B. Dietary Preparation	4
C. Animals	5
D. Animal Maintenance	6
E. Selection of Initial Concentrations	9
F. Experimental Design	9
G. Clinical and Histopathologic Examinations	12
H. Data Recording and Statistical Analyses	14
III. CHRONIC TESTING RESULTS: RATS	19
A. Body Weights and Clinical Observations	19
B. Survival	19
C. Pathology	22
D. Statistical Analyses of Results	26
IV. CHRONIC TESTING RESULTS: MICE	42
A. Body Weights and Clinical Observations	42
B. Survival	42
C. Pathology	45
D. Statistical Analyses of Results	47
V. DISCUSSION	57
VI. BIBLIOGRAPHY	60
APPENDIX A      SUMMARY OF THE INCIDENCE OF NEOPLASMS IN RATS TREATED WITH p-CRESIDINE	A-1
APPENDIX B      SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MICE TREATED WITH p-CRESIDINE	B-1
APPENDIX C      SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN RATS TREATED WITH p-CRESIDINE	C-1
APPENDIX D      SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MICE TREATED WITH p-CRESIDINE	D-1

## LIST OF ILLUSTRATIONS

<u>Figure Number</u>		<u>Page</u>
1	CHEMICAL STRUCTURE OF p-CRESIDINE	2
2	GROWTH CURVES FOR p-CRESIDINE CHRONIC STUDY RATS	20
3	SURVIVAL COMPARISONS OF p-CRESIDINE CHRONIC STUDY RATS	21
4	GROWTH CURVES FOR p-CRESIDINE CHRONIC STUDY MICE	43
5	SURVIVAL COMPARISONS OF p-CRESIDINE CHRONIC STUDY MICE	44

## LIST OF TABLES

<u>Table Number</u>		<u>Page</u>
1	DESIGN SUMMARY FOR FISCHER 344 RATS--p-CRESIDINE FEEDING EXPERIMENT	10
2	DESIGN SUMMARY FOR B6C3F1 MICE--p-CRESIDINE FEEDING EXPERIMENT	11
3	ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT SPECIFIC SITES IN MALE RATS TREATED WITH p-CRESIDINE	27
4	ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT SPECIFIC SITES IN FEMALE RATS TREATED WITH p-CRESIDINE	34
5	TIME-ADJUSTED ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT SPECIFIC SITES IN MALE MICE TREATED WITH p-CRESIDINE	48
6	TIME-ADJUSTED ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT SPECIFIC SITES IN FEMALE MICE TREATED WITH p-CRESIDINE	51

# LIST OF TABLES (Concluded)

<u>Table Number</u>		<u>Page</u>
A1	SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE RATS TREATED WITH p-CRESIDINE	A-3
A2	SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE RATS TREATED WITH p-CRESIDINE	A-8
B1	SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE MICE TREATED WITH p-CRESIDINE	B-3
B2	SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE MICE TREATED WITH p-CRESIDINE	B-7
C1	SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE RATS TREATED WITH p-CRE- SIDINE	C-3
C2	SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE RATS TREATED WITH p-CRE- SIDINE	C-8
D1	SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE MICE TREATED WITH p-CRESI- DINE	D-3
D2	SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE MICE TREATED WITH p-CRE- SIDINE	D-6





## I. INTRODUCTION

p-Cresidine (Figure 1) (NCI No. C02982), used in the production of various azo dyes, was selected for bioassay by the National Cancer Institute in response to the high incidence of bladder cancer observed among dye manufacturing industry workers (Anthony and Thomas, 1970; Wynder et al., 1963). Aromatic amines are one class of chemicals believed to contribute to the increased cancer risk in this industry (Wynder et al., 1963).

The Chemical Abstracts Service (CAS) Ninth Collective Index (1977) name for this compound is 2-methoxy-5-methylbenzenamine.\* It is also known as 2-methoxy-5-methylaniline; 5-methyl-o-anisidine; m-amino-p-cresol methyl ether; MASO; and simply as cresidine.


p-Cresidine appears to be used solely as an intermediate for dyes and pigments. It is believed to be used in the preparation of eleven dyes which are produced commercially in the United States: C.I. (Colour Index) Direct Black 17, Direct Blue 67, Direct Blue 126, Direct Green 26, Direct Orange 34, Direct Orange 83, Direct Red 79, Direct Violet 9, Direct Violet 51, Direct Yellow 41, and Disperse Black 2 (Society of Dyers and Colourists, 1971; as cited in Urso, 1977).

Precise production figures for p-cresidine are not available; however, the Sherwin-Williams Company, which produces p-cresidine,

---

\*The CAS registry number is 120-71-8.





**FIGURE 1**  
**CHEMICAL STRUCTURE OF p-CRESIDINE**

estimates that average use of the compound during the period 1971 to 1976 was about 590,000 pounds annually (Compton, 1977).

Since p-cresidine is used solely as an intermediate in dye manufacture, potential for exposure is primarily limited to workers in dye production facilities.

While adequate chronic toxicity data for p-cresidine are not available, a study of workmen involved in the manufacture and handling of the compound, conducted by the Sherwin-Williams Company, revealed no adverse health effects related to p-cresidine exposure (Compton, 1977).

## II. MATERIALS AND METHODS

### A. Chemicals

p-Cresidine was purchased from Carroll Products, Wood River Junction, Rhode Island by the NCI for Mason Research Institute, Worcester, Massachusetts and chemical analysis was performed by Midwest Research Institute, Kansas City, Missouri. The experimentally determined melting point of 51.5° to 52.5°C, due to its narrow range and proximity to the literature value of 51° to 52°C (Pollock and Stevens, 1965), suggested a compound of high purity. Thin-layer chromatography using a benzene:ethyl acetate solvent system and visualized with ultraviolet light indicated the presence of three impurities, two less motile and one more motile than the major compound. Infrared analysis was consistent with the structure of the compound. Ultraviolet analysis indicated  $\lambda_{\max}$  at 290 and 237 nm with respective molar extinction coefficients ( $\epsilon$ ) of 3312 and 6820. The literature values are  $\lambda_{\max}$  = 290 and 236 nm with respective  $\epsilon$  values of 3620 and 7120 (Mehta, 1978). Titration of the amine function with perchloric acid indicated a purity of 98.9  $\pm$  0.1 percent.

Throughout this report the term p-cresidine is used to represent this technical-grade material.

### B. Dietary Preparation

The basal laboratory diet for both dosed and control animals consisted of Wayne Lab-Blox<sup>®</sup> (Allied Mills, Inc., Chicago, Illinois).

p-Cresidine was administered to the dosed animals as a component of the diet.

The brown amorphous chemical was removed from its shipping container and stored in an amber glass stock bottle inside a plastic bucket. The p-cresidine was weighed out in proper amounts from the stock bottle under an exhaust hood. The compound was dissolved with an appropriate amount of acetone (0.016 percent by volume of the final mix) with the help of a mortar and pestle. This liquid mixture was hand-blended in an aluminum bowl with an aliquot of the ground feed until all liquid was absorbed and visual homogeneity was attained. This premix was then placed into a 6 kg capacity Patterson-Kelley standard model twin-shell stainless steel V-blender along with the remainder of the diet. The blender was sealed and operated for 20 minutes. The mixture was then emptied into stainless steel trays and allowed to sit under the exhaust hood for 2 and 1/2 hours while being stirred with a glass rod every 1/2 hour, allowing evaporation of a large proportion of the acetone.

Prepared diets were placed in double clear plastic bags and stored in the dark at 4°C. Diets were prepared weekly and any unused portions were discarded after 2 weeks.

#### C. Animals

Two animal species, rats and mice, were used in the carcinogenicity bioassay. Fischer 344 rats and B6C3F1 mice were obtained through

contracts of the Division of Cancer Treatment, National Cancer Institute from Charles River Breeding Laboratories, Wilmington, Massachusetts. Dosed and control animals for both species were received in separate shipments.

Upon arrival a sample of animals was examined for parasites and other signs of disease. Animals to be used in the chronic study were quarantined by species for 2 weeks prior to initiation of test. Animals were assigned to groups and distributed among cages so that average body weight per cage was approximately equal for a given species and sex.

#### D. Animal Maintenance

All animals were housed by species in rooms having a temperature range of 23° to 34°C. Incoming air was filtered through Tri-Dek® 15/40 denier Dacron® filters (Tri-Dim Filter Corp., Hawthorne, New Jersey) providing six changes of room air per hour. Fluorescent lighting was provided on a 12-hour-daily cycle.

Rats were housed five per cage by sex. During quarantine and for the first 12 months of study, dosed rats were housed in galvanized steel wire-mesh cages (Fenco Cage Products, Boston, Massachusetts) suspended over newspapers. Control rats were held in galvanized cages for the first 14 months of study. Newspapers under cages were replaced daily and cages and racks washed weekly. For the remainder of the study, all rats were housed in suspended polycarbonate cages (Lab Products, Inc., Garfield, New Jersey) equipped with disposable



nonwoven fiber filter sheets. Clean bedding and cages were provided twice weekly. SAN-I-CEL® corncob bedding (Paxton Processing Company, Paxton, Illinois) was used during the first 6 months of polycarbonate caging. Aspen hardwood chip bedding (American Excelsior Company, Baltimore, Maryland) was used for the remainder of the study. Stainless steel cage racks (Fenco Cage Products) were cleaned once every 2 weeks, and disposable filters were replaced at that time.

Mice were housed by sex in polycarbonate cages fitted with perforated stainless steel lids (Lab Products, Inc.). Nonwoven fiber filter bonnets were used over cage lids. Dosed mice were housed ten per cage for the first 9 months and five per cage thereafter. Control mice were changed from ten per cage to five per cage after 11 months. Clean cages, lids, and bedding were provided three times per week when cage populations were ten and twice per week when cage populations were five. SAN-I-CEL® was used for the first 12 months for control mice, and for the first 10 months for dosed mice. Bed-o-Cobs® corncob bedding (The Andersons Cob Division, Maumee, Ohio) was used for 8 months subsequent to utilization of SAN-I-CEL®. Aspen bedding was then used until completion of the study for all groups of mice. Reusable filter bonnets and pipe racks were sanitized every 2 weeks throughout the study.

Water was available from 250 ml water bottles equipped with rubber stoppers and stainless steel sipper tubes. Bottles were replaced

twice weekly and, for rats only, water was supplied as needed between changes. Food and water were available ad libitum.

During the period of chemical administration, animals were supplied Wayne Lab-Blox<sup>®</sup> containing the appropriate concentrations of p-cresidine and acetone. The basal laboratory diet was supplied to control animals during the same period. Meal was distributed from Alpine<sup>®</sup> aluminum feed cups (Curtin Matheson Scientific, Inc., Woburn, Massachusetts) to rats while in galvanized-steel cages. While in polycarbonate cages, rats were fed from stainless steel gangstyle hoppers (Scientific Cages, Inc., Bryan, Texas). Mice were fed from Alpine<sup>®</sup> feed cups for the first 12 months (dosed groups) or 14 months (controls) of study. Stainless steel gangstyle feed hoppers were used for the remainder of the test. Food hoppers were changed on the same schedule as were cages. Food was replenished daily in Alpine<sup>®</sup> feed cups.

Dosed and control rats were housed in a room with other rats receiving diets containing\* acetylaminofluorene (53-96-3); 2,3,5,6-tetrachloro-4-nitroanisoie (2438-88-2); 4-chloro-o-phenylenediamine (95-83-0); 1H-benzotriazole (95-14-7); and 4-chloro-m-phenylenediamine (5131-60-2).

Dosed and control mice were housed in a room with other mice receiving diets containing 2-methyl-1-nitroanthraquinone (129-15-7); aniline hydrochloride (142-04-1); acetylaminofluorene (53-96-3); fenaminosulf (140-56-7); and 4-chloro-m-phenylenediamine (5131-60-2).

---

\*CAS registry numbers are given in parentheses.



#### E. Selection of Initial Concentrations

In order to establish the maximum tolerated concentrations of p-cresidine for administration to dosed animals in the chronic studies, subchronic toxicity tests were conducted with both rats and mice.

Animals of each species were distributed among three groups, each consisting of five males and five females. p-Cresidine was administered to two of the three groups of each species at dietary concentrations of 1.0 and 3.0 percent for 8 weeks. The third group of each species served as a control group, receiving only the basal laboratory diet.

As deaths occurred among all groups except male rats receiving 3.0 percent p-cresidine in the feed, the high concentration chosen for the chronic study was 1.0 percent p-cresidine for both rats and mice.

#### F. Experimental Design

The experimental design parameters for the chronic study (species, sex, group size, concentrations administered, duration of treated and untreated observation periods, and time-weighted average concentrations) are summarized in Tables 1 and 2.

All animals were approximately 6 weeks old at the time they were placed on test. Dosed rats and mice shared the same median date of birth, while controls were born approximately 7 weeks earlier and placed on test 7 weeks before dosed animals.

TABLE 1

DESIGN SUMMARY FOR FISCHER 344 RATS  
p-CRESIDINE FEEDING EXPERIMENT

	INITIAL GROUP SIZE	p-CRESIDINE CONCENTRATION (PERCENT)	OBSERVATION PERIOD	
			TREATED (WEEKS)	UNTREATED (WEEKS)
<u>MALE</u>				
CONTROL	50	0	0	106
LOW DOSE	50	0.5 0	104	1
HIGH DOSE	50	1.0 0	104	1
<u>FEMALE</u>				
CONTROL	50	0	0	106
LOW DOSE	50	0.5 0	104	2
HIGH DOSE	50	1.0 0	104	2

TABLE 2

DESIGN SUMMARY FOR B6C3F1 MICE  
p-CRESIDINE FEEDING EXPERIMENT

	INITIAL GROUP SIZE	p-CRESIDINE CONCENTRATION (PERCENT)	OBSERVATION PERIOD		TIME-WEIGHTED AVERAGE CONCENTRATION <sup>a</sup>
			TREATED (WEEKS)	UNTREATED (WEEKS)	
<u>MALE</u>					
CONTROL	50	0	0	97	0
<hr/>					
LOW DOSE	50	0.5 0.15 0	21 83	2	0.22
<hr/>					
HIGH DOSE <sup>b</sup>	50	1.0 0.3	21 71		0.46
<hr/> <hr/>					
<u>FEMALE</u>					
CONTROL	50	0	0	97	0
<hr/>					
LOW DOSE	50	0.5 0.15 0	21 83	2	0.22
<hr/>					
HIGH DOSE	50	1.0 0.3 0	21 83	2	0.44

<sup>a</sup> Time-weighted average concentration =  $\frac{\sum(\text{concentration} \times \text{weeks received})}{\sum(\text{weeks receiving chemical})}$

<sup>b</sup> All animals in this group were dead by the end of week 92.

When it was noted that a trace acetone odor remained in the dosed feed, acetone controls were placed on test in addition to the untreated controls. However, these acetone controls were not entirely comparable to dosed animals because one set of acetone controls had an inadequate number of animals (only 10 animals of each sex and species) and were treated for parasites with piperazine adipate. A second set of acetone controls was not utilized in comparisons with dosed animals because they were obtained from a different supplier than the other groups.

The initial concentrations of p-cresidine used for both rats and mice were 1.0 and 0.5 percent. Throughout this report those animals initially receiving the former concentration are referred to as the high dose groups, while those initially receiving the latter concentration are referred to as the low dose groups. These concentrations were utilized throughout the study for rats; however, after 21 weeks, the concentrations administered to mice were decreased to 0.3 and 0.15 percent for the high and low dose groups, respectively. p-Cresidine was administered in the diet to dosed animals for a period of 104 weeks (except for high dose male mice which were all dead by week 92), followed by an observation period of up to 2 weeks.

For both species, control animals had the basal laboratory diet available ad libitum during the bioassay.

#### G. Clinical and Histopathologic Examinations

Animals were weighed immediately prior to initiation of the experiment. Body weights were recorded twice weekly for the first 12



weeks of the study and at monthly intervals thereafter. From the first day, all animals were inspected twice daily for mortality. Food consumption, for two cages from each group, was monitored for seven consecutive days once a month for the first nine months of the bioassay and for three consecutive days each month thereafter. The presence of tissue masses and lesions was determined by monthly observation and palpation of each animal.

A necropsy was performed on each animal regardless of whether it died, was killed when moribund, or was sacrificed at the end of the bioassay. The animals were euthanized by carbon dioxide inhalation, and were immediately necropsied. The histopathologic examination consisted of gross and microscopic examination of major tissues, organs, and gross lesions taken from sacrificed animals and, whenever possible, from animals found dead.

Tissues were preserved in 10 percent buffered formalin, embedded in paraffin, sectioned, and stained with hematoxylin and eosin prior to microscopic examination. An occasional section was subjected to special staining techniques for more definitive diagnosis.

Slides were prepared from the following tissues: skin, subcutaneous tissue, lungs and bronchi, trachea, nasal cavity, bone marrow, spleen, lymph nodes, thymus, heart, salivary gland, liver, gallbladder (mice), pancreas, esophagus, stomach, small intestine, large intestine, kidney, urinary bladder, seminal vesicle, pituitary, adrenal, thyroid, parathyroid, testis, prostate, brain, uterus, mammary gland, and ovary.

A few tissues were not examined for some animals, particularly for those that died early. Also, some animals were missing, cannibalized, or judged to be in such an advanced state of autolysis as to preclude histopathologic interpretation. Thus, the number of animals for which particular organs, tissues, or lesions were examined microscopically varies and does not necessarily represent the number of animals that were placed on experiment in each group.

#### H. Data Recording and Statistical Analyses

Pertinent data on this experiment have been recorded in an automatic data processing system, the Carcinogenesis Bioassay Data System (Linhart et al., 1974). The data elements include descriptive information on the chemicals, animals, experimental design, clinical observations, survival, body weight, and individual pathologic results, as recommended by the International Union Against Cancer (Berenblum, 1969). Data tables were generated for verification of data transcription and for statistical review.

These data were analyzed using the statistical techniques described in this section. Those analyses of the experimental results that bear on the possibility of carcinogenicity are discussed in the statistical narrative sections.

Probabilities of survival were estimated by the product-limit procedure of Kaplan and Meier (1958) and are presented in this report in the form of graphs. Animals were statistically censored as of the time that they died of other than natural causes or were found to be

missing; animals dying from natural causes were not statistically censored. Statistical analyses for a possible dose-related effect on survival used the method of Cox (1972) when testing two groups for equality and used Tarone's (1975) extensions of Cox's methods when testing a dose-related trend. One-tailed P-values have been reported for all tests except the departure from linearity test, which is only reported when its two-tailed P-value is less than 0.05.

The incidence of neoplastic or nonneoplastic lesions has been given as the ratio of the number of animals bearing such lesions at a specific anatomic site (numerator) to the number of animals in which that site was examined (denominator). In most instances, the denominators included only those animals for which that site was examined histologically. However, when macroscopic examination was required to detect lesions prior to histologic sampling (e.g., skin or mammary tumors), or when lesions could have appeared at multiple sites (e.g., lymphomas), the denominators consist of the numbers of animals necropsied.

The purpose of the statistical analyses of tumor incidence is to determine whether animals receiving the test chemical developed a significantly higher proportion of tumors than did the control animals. As a part of these analyses, the one-tailed Fisher exact test (Cox, 1970, pp. 48-52) was used to compare the tumor incidence of a control group to that of a group of treated animals at each dose level. When results for a number of treated groups,  $k$ , are compared simultaneously



with those for a control group, a correction to ensure an overall significance level of 0.05 may be made. The Bonferroni inequality (Miller, 1966, pp. 6-10) requires that the P-value for any comparison be less than or equal to  $0.05/k$ . In cases where this correction was used, it is discussed in the narrative section. It is not, however, presented in the tables, where the Fisher exact P-values are shown.

The Cochran-Armitage test for linear trend in proportions, with continuity correction (Armitage, 1971, pp. 362-365), was also used when appropriate. Under the assumption of a linear trend, this test determined if the slope of the dose-response curve is different from zero at the one-tailed 0.05 level of significance. Unless otherwise noted, the direction of the significant trend was a positive dose relationship. This method also provides a two-tailed test of departure from linear trend.

A time-adjusted analysis was applied when numerous early deaths resulted from causes that were not associated with the formation of tumors. In this analysis, deaths that occurred before the first tumor was observed were excluded by basing the statistical tests on animals that survived at least 52 weeks, unless a tumor was found at the anatomic site of interest before week 52. When such an early tumor was found, comparisons were based exclusively on animals that survived at least as long as the animal in which the first tumor was found. Once this reduced set of data was obtained, the standard

procedures for analyses of the incidence of tumors (Fisher exact tests, Cochran-Armitage tests, etc.) were followed.

When appropriate, life-table methods were used to analyze the incidence of tumors. Curves of the proportions surviving without an observed tumor were computed as in Saffiotti et al. (1972). The week during which animals died naturally or were sacrificed was entered as the time point of tumor observation. Cox's methods of comparing these curves were used for two groups; Tarone's extension to testing for linear trend was used for three groups. The statistical tests for the incidence of tumors which used life-table methods were one-tailed and, unless otherwise noted, in the direction of a positive dose relationship. Significant departures from linearity ( $P < 0.05$ , two-tailed test) were also noted.

The approximate 95 percent confidence interval for the relative risk of each dosed group compared to its control was calculated from the exact interval on the odds ratio (Gart, 1971). The relative risk is defined as  $p_t/p_c$  where  $p_t$  is the true binomial probability of the incidence of a specific type of tumor in a treated group of animals and  $p_c$  is the true probability of the spontaneous incidence of the same type of tumor in a control group. The hypothesis of equality between the true proportion of a specific tumor in a treated group and the proportion in a control group corresponds to a relative risk of unity. Values in excess of unity represent the condition of a larger proportion in the treated group than in the control.

The lower and upper limits of the confidence interval of the relative risk have been included in the tables of statistical analyses. The interpretation of the limits is that in approximately 95 percent of a large number of identical experiments, the true ratio of the risk in a treated group of animals to that in a control group would be within the interval calculated from the experiment. When the lower limit of the confidence interval is greater than one, it can be inferred that a statistically significant result (a  $P < 0.025$  one-tailed test when the control incidence is not zero,  $P < 0.050$  when the control incidence is zero) has occurred. When the lower limit is less than unity but the upper limit is greater than unity, the lower limit indicates the absence of a significant result while the upper limit indicates that there is a theoretical possibility of the induction of tumors by the test chemical which could not be detected under the conditions of this test.

### III. CHRONIC TESTING RESULTS: RATS

#### A. Body Weights and Clinical Observations

Male and female dosed rats had mean group body weights lower than those of their respective controls (Figure 2), especially the weights of the high dose male group which were markedly decreased throughout the study. Fluctuations in the growth curve may be due to mortality; as the size of the group diminishes, the mean body weight may be subject to wide variations.

Subcutaneous masses on various parts of the body were observed in 2 control, 3 low dose, and 1 high dose males and in 3 control, 7 low dose, and 1 high dose females. Cutaneous masses or nodules were detected in 2 control and 3 low dose males and in 2 low dose and 1 high dose females. Skin lesions were observed in 1 control male. Other isolated clinical observations included penile swelling in 1 high dose male, swelling of a single nipple in 1 low dose female, and severe emaciation in 1 low dose male.

#### B. Survival

The estimated probabilities of survival for male and female rats in the control and p-cresidine-dosed groups are shown in Figure 3. For both male and female rats there were significant positive associations between dosage and mortality.

For males five rats from the control group were sacrificed in week 78. Sixty-two percent (31/50) of the high dose, 96 percent







FIGURE 2  
GROWTH CURVES FOR p-CRESIDINE CHRONIC STUDY RATS





FIGURE 3  
SURVIVAL COMPARISONS OF p-CRESIDINE CHRONIC STUDY RATS

(48/50) of the low dose, and 94 percent (47/50) of the control rats survived on test at least 75 weeks.

For female rats five of the controls were sacrificed in week 78. Seventy-six percent (38/50) of the high dose, 98 percent (49/50) of the low dose, and 96 percent (48/50) of the control rats survived on test at least 75 weeks.

#### C. Pathology

Histopathologic findings on neoplasms in rats are summarized in Appendix A (Tables A1 and A2); findings on nonneoplastic lesions are summarized in Appendix C (Tables C1 and C2).

Both male and female rats fed p-cresidine showed compound-related effects in the urinary bladder and in the epithelium of the nasal cavity.

No neoplasms of the bladder were found in the control rats. In contrast, both male and female rats fed p-cresidine developed numerous dose-related neoplasms or hyperplasias in the bladder. The incidences of these lesions are tabulated in the table below:



<u>MALE</u>	<u>Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Number of animals with tissues examined histopathologically	(48)	(48)	(47)
<u>Urinary Bladder</u>			
Hyperplasia	0	11 (23%)	2 (4%)
Undifferentiated Carcinoma	0	0	1 (2%)
Papillary Carcinoma	0	0	4 (9%)
Squamous-Cell Carcinoma	0	1 (2%)	6 (13%)
Transitional-Cell Papilloma	0	14 (29%)	3 (6%)
Transitional-Cell Carcinoma	0	15 (31%)	31 (66%)
Leiomyosarcoma	0	0	2 (4%)
<u>FEMALE</u>			
Number of animals with tissues examined histopathologically	(47)	(49)	(46)
<u>Urinary Bladder</u>			
Hyperplasia	0	11 (22%)	1 (2%)
Neoplasm NOS	0	0	2 (4%)
Papillary Carcinoma	0	0	1 (2%)
Squamous-Cell Carcinoma	0	9 (18%)	7 (15%)
Transitional-Cell Papilloma	0	6 (12%)	2 (4%)
Transitional-Cell Carcinoma	0	22 (45%)	33 (72%)
Leiomyosarcoma	0	2 (4%)	3 (7%)

Hyperplasia was defined as the presence of more than three layers of epithelial cells lining the fully inflated bladder. Nuclei of these cells were histologically normal.

The neoplasms were classified as papillomas when they consisted of histologically benign, regularly arranged, transitional epithelium covering a papillary core composed of fibrovascular tissue. Tumors were diagnosed as transitional-cell carcinomas when nuclei of the transitional cells became pleomorphic and hyperchromatic and mitotic figures became frequent. They were classified as squamous-cell

carcinomas when, in addition to nuclear criteria of malignancy, the tumor cells showed cytoplasmic keratinization or developed intercellular bridges. There were no sharp boundary lines between these categories which formed a continuous spectrum of progressively malignant change. The carcinomas were invasive and a few of these tumors metastasized to distant organs. A few rats fed p-cresidine developed leiomyosarcomas of the bladder wall which tended to occur in association with malignant tumors of the bladder epithelium. There were neoplasms of the transitional-cell epithelium of the renal pelvis in 1/49 low dose female rats and in 2/49 low dose males. One high dose female developed an undifferentiated carcinoma of the renal parenchyma.

No nasopharyngeal neoplasms were found in the control groups. The incidence of neoplasms observed in the nasal cavity is summarized in the following table:

<u>MALES</u>	<u>Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Number of animals necropsied	(48)	(50)	(47)
<u>Nasal Cavity</u>			
Neoplasm NOS, Malignant	0	0	1(2%)
Carcinoma NOS	0	0	1(2%)
Adenoma NOS	0	1(2%)	0
Neuroblastoma	0	0	1(2%)
Olfactory Neuroblastoma	0	1(2%)	20(43%)

<u>FEMALES</u>	<u>Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Number of animals necropsied	(50)	(50)	(49)
<u>Nasal Cavity</u>			
Squamous-Cell Carcinoma	0	0	1(2%)
Squamous-Cell Carcinoma, Invasive	0	0	0
Adenocarcinoma NOS	0	0	1(2%)
Olfactory Neuroblastoma	0	0	11(22%)

The rats fed p-cresidine developed dose-related olfactory neuroblastomas. These tumors consisted of masses of cells having scant cytoplasm and large, ovoid nuclei with prominent chromatin. They were divided into lobules by very fine fibrovascular septae. In many tumors, rosettes, pseudorosettes, or palisade arrangement of cells suggested a neural origin. They frequently exhibited osteoclastic activity and local invasion, destroying the cribriform plate of the skull and invading the forebrain. Some tumors showed focal hemorrhagic necrosis. In addition, there was an adenoma, an adenocarcinoma, a squamous-cell carcinoma, and a polyp involving the nasal cavity of dosed rats. Males appeared to be more susceptible to nasal tumors than females.

p-Cresidine feeding had an equivocal effect on the liver of the rats. Males receiving the low dose of the compound developed several (10/49 [20 percent]) neoplastic nodules plus a few hepatic carcinomas. However, neoplastic nodules were seldom seen in high dose males (1/46 [2 percent]) and low dose females (4/48 [8 percent]) and were not seen at all in high dose females. Only a few carcinomas were observed in

dosed male rats. The criteria used for the diagnosis of these lesions were those of Squire and Levitt (1975).

No tumors of the preputial or clitoral glands were seen in controls; 3/50 (6 percent) occurred in low dose males, 3/50 (6 percent) in low dose females, and 2/47 (4 percent) in high dose males. There were gliomas in 2/48 (4 percent) low dose females, and in 3/43 (7 percent) high dose males. As the high dose of p-cresidine shortened the survival of both sexes by inducing massive bladder tumors, an effect on other organ systems may have been obscured.

Based on the results of this pathologic examination, p-cresidine was considered to be carcinogenic in Fischer 344 rats because it induced neoplasms of the urinary bladder and olfactory neuroblastomas of the nasal cavity. Neoplastic nodules of the liver may have been induced in the male rat.

#### D. Statistical Analyses of Results

The results of the statistical analyses of tumor incidence in rats are summarized in Tables 3 and 4. The analysis is included for every type of malignant tumor in either sex where at least two such tumors were observed in at least one of the control or p-cresidine-dosed groups and where such tumors were observed in at least 5 percent of the group.

In both male and female rats, the combined incidence of undifferentiated carcinomas, papillary carcinomas, squamous-cell carcinomas, transitional-cell papillomas, or transitional-cell carcinomas of the



ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT  
SPECIFIC SITES IN MALE RATS TREATED WITH P-CRESIDINE<sup>a</sup>

TOPOGRAPHY:MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Subcutaneous Tissue: Fibroma <sup>b</sup>	3/48(0.06)	2/50(0.04)	1/47(0.02)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	0.640	0.340
Lower Limit	---	0.055	0.007
Upper Limit	---	5.345	4.058
Weeks to First Observed Tumor	105	105	105
Nasal Cavity and Nasopharynx: Malignant Neoplasm NOS; Carcinoma NOS; or Adenocarcinoma NOS <sup>b</sup>	0/48(0.00)	0/50(0.00)	3/47(0.06)
P Values <sup>c</sup>	P = 0.035	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	---	Infinite
Lower Limit	---	---	0.615
Upper Limit	---	---	Infinite
Weeks to First Observed Tumor	---	---	67
Nasal Cavity: Olfactory Neuroblastoma or Neuroblastoma <sup>b</sup>	0/48(0.00)	1/50(0.02)	21/47(0.45)
P Values <sup>c</sup>	P < 0.001	N.S.	P < 0.001
Departure from Linear Trend <sup>e</sup>	P = 0.001	---	---
Relative Risk (Control) <sup>d</sup>	---	Infinite	Infinite
Lower Limit	---	0.051	6.974
Upper Limit	---	Infinite	Infinite
Weeks to First Observed Tumor	---	84	48



TABLE 3 (Continued)

TOPOGRAPHY: MORPHOLOGY		CONTROL	LOW DOSE	HIGH DOSE
Nasal Cavity: Neoplasm NOS, Malignant; Carcinoma NOS; Adenoma NOS; Neuroblastoma; or Olfactory Neuroblastoma <sup>b</sup>		0/48(0.00)	2/50(0.04)	23/47(0.49)
P Values <sup>c</sup>		P < 0.001	N.S.	P < 0.001
Departure from Linear Trend <sup>e</sup>		P = 0.002	---	---
Relative Risk (Control) <sup>d</sup>		---	Infinitive	Infinitive
Lower Limit		---	0.284	7.699
Upper Limit		---	Infinitive	Infinitive
Weeks to First Observed Tumor		---	84	48
Lung: Alveolar/Bronchiolar Adenoma <sup>b</sup>		3/48(0.06)	0/49(0.00)	0/45(0.00)
P Values <sup>c</sup>		P = 0.040(N)	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>		---	0.000	0.000
Lower Limit		---	0.000	0.000
Upper Limit		---	1.628	1.768
Weeks to First Observed Tumor		105	---	---
Hematopoietic System: Leukemia or Malignant Lymphoma <sup>b</sup>		7/48(0.15)	1/50(0.02)	2/47(0.04)
P Values <sup>c</sup>		P = 0.037(N)	P = 0.026(N)	N.S.
Relative Risk (Control) <sup>d</sup>		---	0.137	0.292
Lower Limit		---	0.003	0.031
Upper Limit		---	1.009	1.436
Weeks to First Observed Tumor		80	105	72

TABLE 3 (Continued)

TOPOGRAPHY: MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Liver: Hepatocellular Carcinoma, or Mixed Hepato/Cholangio Carcinoma <sup>b</sup>	0/48(0.00)	3/49(0.06)	1/46(0.02)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	Infinite	Infinite
Lower Limit	---	0.590	0.056
Upper Limit	---	Infinite	Infinite
Weeks to First Observed Tumor	---	105	78
Liver: Neoplastic Nodule, Hepatocellular Carcinoma, or Mixed Hepato/Cholangio Carcinoma <sup>b</sup>	0/48(0.00)	13/49(0.27)	2/46(0.04)
P Values <sup>c</sup>	N.S.	P < 0.001	N.S.
Departure from Linear Trend <sup>e</sup>	P < 0.001	---	---
Relative Risk (Control) <sup>d</sup>	---	Infinite	Infinite
Lower Limit	---	3.938	0.309
Upper Limit	---	Infinite	Infinite
Weeks to First Observed Tumor	---	105	78

TABLE 3 (Continued)

TOPOGRAPHY: MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Urinary Bladder: Undifferentiated Carcinoma, Papillary Carcinoma, Squamous-Cell Carcinoma, Transitional-Cell Papilloma, or Transitional-Cell Carcinoma <sup>b</sup>	0/48(0.00)	30/48(0.63)	44/47(0.94)
P Values <sup>c</sup>	P < 0.001	P < 0.001	P < 0.001
Relative Risk (Control) <sup>d</sup>	---	Infinitive	Infinitive
Lower Limit	---	10.044	16.426
Upper Limit	---	Infinitive	Infinitive
Weeks to First Observed Tumor	---	93	51
Pituitary: Adenoma NOS, Chromophobe Adenoma, or Basophil Adenoma <sup>b</sup>	10/45(0.22)	9/48(0.19)	3/38(0.08)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	0.844	0.355
Lower Limit	---	0.335	0.067
Upper Limit	---	2.095	1.260
Weeks to First Observed Tumor	103	85	56
Adrenal: Cortical Adenoma <sup>b</sup>	0/46(0.00)	3/48(0.06)	0.44(0.00)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	Infinitive	---
Lower Limit	---	0.578	---
Upper Limit	---	Infinitive	---
Weeks to First Observed Tumor	---	102	---

TABLE 3 (Continued)

TOPOGRAPHY: MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Adrenal: Pheochromocytoma <sup>b</sup>	4/46(0.09)	7/48(0.15)	3/44(0.07)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	1.677	0.784
Lower Limit	---	0.459	0.121
Upper Limit	---	7.336	4.367
Weeks to First Observed Tumor	78	102	80
Thyroid: C-Cell Carcinoma, or C-Cell Adenoma <sup>b</sup>	5/43(0.12)	2/41(0.05)	2/40(0.05)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	0.420	0.430
Lower Limit	---	0.042	0.043
Upper Limit	---	2.398	2.455
Weeks to First Observed Tumor	95	105	93
Pancreatic Islets: Islet-Cell Adenoma, or Islet-Cell Carcinoma <sup>b</sup>	0/44(0.00)	2/48(0.04)	2/40(0.05)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	Infinite	Infinite
Lower Limit	---	0.272	0.327
Upper Limit	---	Infinite	Infinite
Weeks to First Observed Tumor	---	105	97

TABLE 3 (Continued)

TOPOGRAPHY:MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Preputial Gland: Adenoma NOS, or Carcinomas NOS <sup>b</sup>	0/48(0.00)	3/50(0.06)	1/47(0.02)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	Infinitive	Infinitive
Lower Limit	---	0.578	0.055
Upper Limit	---	Infinitive	Infinitive
Weeks to First Observed Tumor	---	79	97
Testis: Interstitial-Cell Tumor <sup>b</sup>	37/48(0.77)	45/47(0.96)	23/46(0.50)
P Values <sup>c</sup>	P = 0.003(N)	P = 0.008	P = 0.006(N)
Departure from Linear Trend <sup>e</sup>	P < 0.001	---	---
Relative Risk (Control) <sup>d</sup>	---	1.242	0.649
Lower Limit	---	1.038	0.471
Upper Limit	---	1.354	0.916
Weeks to First Observed Tumor	78	79	52
Brain and Brain Stem: Glioma NOS, or Oligodendroglioma <sup>b</sup>	0/46(0.00)	0/48(0.00)	3/43(0.07)
P Values <sup>c</sup>	P = 0.032	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	---	Infinitive
Lower Limit	---	---	0.645
Upper Limit	---	---	Infinitive
Weeks to First Observed Tumor	---	---	56



TABLE 3 (Concluded)

<sup>a</sup>Treated groups received doses of 0.5 or 1.0 percent in feed.

<sup>b</sup>Number of tumor-bearing animals/number of animals examined at site (proportion).

<sup>c</sup>The probability level for the Cochran-Armitage test is given beneath the incidence of tumors in the control group when  $P < 0.05$ ; otherwise, not significant (N.S.) is indicated. The probability level for the Fisher exact test for the comparison of a treated group with the control group is given beneath the incidence of tumors in the treated group when  $P < 0.05$ ; otherwise, not significant (N.S.) is indicated. For both Cochran-Armitage and Fisher exact tests a negative designation (N) indicates a lower incidence in the treated group(s) than in the control group.

<sup>d</sup>The 95 percent confidence interval on the relative risk of the treated group to the control group.

<sup>e</sup>The probability level of the test for departure from linear trend is given beneath the control group when  $P < 0.05$ .

TABLE 4

ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT  
SPECIFIC SITES IN FEMALE RATS TREATED WITH P-CRESIDINE<sup>a</sup>

TOPOGRAPHY:MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Nasal Cavity: Olfactory Neuroblastoma <sup>b</sup>	0/50(0.00)	0/50(0.00)	11/49(0.22)
P Values <sup>c</sup>	P < 0.001	N.S.	P < 0.001
Departure from Linear Trend <sup>e</sup>	P = 0.013	---	---
Relative Risk (Control) <sup>d</sup>	---	---	Infinite
Lower Limit	---	---	3.389
Upper Limit	---	---	Infinite
Weeks to First Observed Tumor	---	---	62
Hematopoietic System: Leukemia or Malignant Lymphoma <sup>b</sup>			
	4/50(0.08)	0/50(0.00)	0/49(0.00)
P Values <sup>c</sup>	P = 0.016(N)	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	0.000	0.000
Lower Limit	---	0.000	0.000
Upper Limit	---	1.079	1.100
Weeks to First Observed Tumor	102	---	---
Liver: Neoplastic Nodule <sup>b</sup>			
	0/50(0.00)	4/48(0.08)	0/48(0.00)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	Infinite	---
Lower Limit	---	0.966	---
Upper Limit	---	Infinite	---
Weeks to First Observed Tumor	---	76	---

TABLE 4 (Continued)

TOPOGRAPHY:MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Urinary Bladder: Leiomyosarcoma <sup>b</sup>	0/47(0.00)	2/49(0.04)	3/46(0.07)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	Infinit	Infinit
Lower Limit	---	0.284	0.616
Upper Limit	---	Infinit	Infinit
Weeks to First Observed Tumor	---	106	95
Urinary Bladder: Neoplasm NOS, Papillary Carcinoma, Squamous- Cell Carcinoma, or Transitional- Cell Carcinoma <sup>b</sup>	0/47(0.00)	31/49(0.63)	43/46(0.93)
P Values <sup>c</sup>	P < 0.001	P < 0.001	P < 0.001
Departure from Linear Trend <sup>e</sup>	P = 0.043	---	---
Relative Risk (Control) <sup>d</sup>	---	Infinit	Infinit
Lower Limit	---	9.978	16.054
Upper Limit	---	Infinit	Infinit
Weeks to First Observed Tumor	---	88	59

TABLE 4 (Continued)

TOPOGRAPHY:MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Pituitary: Adenoma NOS, or Chromophobe Adenomab	17/40(0.43)	11/45(0.24)	7/40(0.18)
P Values <sup>c</sup>	P = 0.009(N)	N.S.	P = 0.013(N)
Relative Risk (Control) <sup>d</sup>	---	0.575	0.412
Lower Limit	---	0.282	0.165
Upper Limit	---	1.138	0.918
Weeks to First Observed Tumor	101	79	70
Adrenal: Cortical Adenomab	0/48(0.00)	5/48(0.10)	1/44(0.02)
P Values <sup>c</sup>	N.S.	P = 0.028	N.S.
Departure from Linear Trend <sup>e</sup>	P = 0.010	---	---
Relative Risk (Control) <sup>d</sup>	---	Infinite	Infinite
Lower Limit	---	1.263	0.059
Upper Limit	---	Infinite	Infinite
Weeks to First Observed Tumor	---	79	86
Thyroid: Follicular-Cell Carcinomab	0/43(0.00)	2/42(0.05)	0/40(0.00)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	Infinite	---
Lower Limit	---	0.304	---
Upper Limit	---	Infinite	---
Weeks to First Observed Tumor	---	106	---

TABLE 4 (Continued)

TOPOGRAPHY: MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Thyroid: C-Cell Adenoma <sup>b</sup>	0/43(0.00)	4/42(0.10)	0/40(0.00)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	Infinite	---
Lower Limit	---	0.954	---
Upper Limit	---	Infinite	---
Weeks to First Observed Tumor	---	103	---
Mammary Gland: Fibroadenoma <sup>b</sup>	6/50(0.12)	4/50(0.08)	1/49(0.02)
P Values <sup>c</sup>	P = 0.045(N)	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	0.667	0.170
Lower Limit	---	0.147	0.004
Upper Limit	---	2.635	1.328
Weeks to First Observed Tumor	106	106	79
Uterus: Endometrial Stromal Polyp <sup>b</sup>	2/48(0.04)	8/47(0.17)	4/45(0.09)
P Values <sup>c</sup>	N.S.	P = 0.042	N.S.
Departure from Linear Trend <sup>e</sup>	P = 0.050	---	---
Relative Risk (Control) <sup>d</sup>	---	4.085	2.133
Lower Limit	---	0.872	0.322
Upper Limit	---	37.831	22.646
Weeks to First Observed Tumor	106	80	82



TABLE 4 (Concluded)

a	Treated groups received doses of 0.5 or 1.0 in feed.
b	Number of tumor-bearing animals/number of animals examined at site (proportion).
c	The probability level for the Cochran-Armitage test is given beneath the incidence of tumors in the control group when $P < 0.05$ ; otherwise, not significant (N.S.) is indicated. The probability level for the Fisher exact test for the comparison of a treated group with the control group is given beneath the incidence of tumors in the treated group when $P < 0.05$ ; otherwise, not significant (N.S.) is indicated. For both Cochran-Armitage and Fisher exact tests a negative designation (N) indicates a lower incidence in the treated group(s) than in the control group.
d	The 95 percent confidence interval on the relative risk of the treated group to the control group.
e	The probability level of the test for departure from linear trend is given beneath the control group when $P < 0.05$ .

urinary bladder was increased in both dosed groups compared to the control group--appearing as early as week 51 in the males and week 59 in the females. For both sexes, the Cochran-Armitage test indicated a significant ( $P < 0.001$ ) positive association between dose and incidence. For both sexes this was supported by significant ( $P < 0.001$ ) Fisher exact test results for both dose levels compared to the control. Based on these results the administration of p-cresidine was associated with the increased incidence of carcinomas of the urinary bladder in both male and female rats.

Both the incidence of olfactory neuroblastomas in females and the combined incidence of neuroblastomas or olfactory neuroblastomas in males were increased in the high dose groups compared to the controls--appearing as early as week 48 in the males and week 62 in the females. For both sexes the Cochran-Armitage test indicated a significant ( $P < 0.001$ ) positive association between dose and incidence. For both sexes these results were supported by significant ( $P < 0.001$ ) Fisher exact tests comparing the high dose group to the control group. For males the combined incidence of nonspecific neoplasms, malignant carcinomas, or adenocarcinomas of the nasal cavity or nasopharynx was significant for the Cochran-Armitage test, but not for the Fisher exact tests. Based on these results, the administration of p-cresidine was associated with the increased incidence of neuroblastomas and olfactory neuroblastomas of the nasal cavity in both male and female rats.

In male rats the combined incidence of neoplastic nodules of the liver, hepatocellular carcinomas, or mixed hepato/cholangio carcinomas was increased in the low dose group compared to the control. The tumors were found after termination of the study in week 105. The Fisher exact test indicated a significantly ( $P < 0.001$ ) higher incidence in the low dose group than in the control. The high dose Fisher exact comparison and the Cochran-Armitage test, however, were not significant, but these tests must be discounted due to the early deaths from cancer in the high dose group. Based on these results, the administration of p-cresidine was probably associated with the increased incidence of liver neoplasms in the male rats.

In male rats an increase in the combined incidence of gliomas and oligodendrogliomas of the brain or brain stem were observed in the high dose group. The Cochran-Armitage test indicated a significant ( $P = 0.032$ ) positive association between dosage and incidence. This was not supported, however, by significant Fisher exact tests.

In males for lung neoplasms and for the incidence of leukemia or malignant lymphomas the Cochran-Armitage test was significant in the negative direction, but the Fisher exact tests were not significant under the Bonferroni criterion. Similarly, in the female rats the results did not impute any real importance to adrenal cortical adenomas, to mammary fibroadenomas, to endometrial stromal polyps, or to leukemias or malignant lymphomas.

The possibility of a negative association was observed between chemical administration and the incidence of pituitary adenomas in

female rats and the incidence of interstitial-cell tumors of the testis in males. This may have been due to the fact that survival was markedly better in the control and low dose groups than in the high dose group.

In summary, the administration of p-cresidine was associated with the incidences of carcinomas of the urinary bladder and of olfactory neuroblastomas in both males and females, and possibly with the incidence of liver neoplasms in males.



#### IV. CHRONIC TESTING RESULTS: MICE

##### A. Body Weights and Clinical Observations

The mean body weights for dosed mice of both sexes were consistently lower than those for the control groups (Figure 4). Fluctuations in the growth curve may be due to mortality; as the size of the group diminishes, the mean body weight may be subject to wide variations.

Only two dosed mice exhibited clinical signs. These were 1 low dose male with a hard mass in the abdominal cavity and 1 low dose female with a swollen, infected eye.

##### B. Survival

The estimated probabilities of survival for male and female mice in the control and p-cresidine-dosed groups are shown in Figure 5. For male mice and for female mice the Tarone tests showed significant positive associations between dosage and mortality.

For males five of the control mice were sacrificed in week 78. High early mortality was observed in the high dose group, with 50 percent (25/50) of the males dead by week 52 and 90 percent (45/50) dead by week 73. With 27 of these 45 mice observed to have a tumor of the urinary bladder, it seemed that mortality was associated with tumor incidence. Fifty percent (25/50) of the low dose and 98 percent (49/50) of the control mice survived on test for at least 75 weeks.







FIGURE 4  
GROWTH CURVES FOR p-CRESIDINE CHRONIC STUDY MICE





FIGURE 5  
SURVIVAL COMPARISONS OF p-CRESIDINE CHRONIC STUDY MICE

For female mice five of the controls were sacrificed in week 78. Twenty-eight percent (14/50) of the high dose, 78 percent (39/50) of the low dose, and 90 percent (45/50) of the control mice survived on test for at least 75 weeks.

### C. Pathology

Histopathologic findings on neoplasms in mice are summarized in Appendix B (Tables B1 and B2); findings on nonneoplastic lesions are summarized in Appendix D (Tables D1 and D2).

p-Cresidine feeding produced a compound-related effect in the urinary bladder. No neoplasms of the urinary bladder occurred in the control mice. In the mice fed p-cresidine, there was a dose-related spectrum of bladder neoplasms which resembled rat bladder neoplasms previously described in this report. The hyperplasias and neoplasms of the urinary bladder in the mice are summarized in the table below:

<u>MALE</u>	<u>Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Number of animals with tissues examined histopathologically	(50)	(42)	(35)
<u>Urinary Bladder</u>			
Hyperplasia	0	0	3(9%)
Dysplasia	0	0	1(3%)
Squamous-Cell Carcinoma	0	27(64%)	28(80%)
Transitional-Cell Carcinoma	0	13(31%)	3(9%)
Leiomyosarcoma	0	2(5%)	4(11%)

## FEMALE

Number of animals with tissues examined histopathologically	(45)	(47)	(48)
<u>Urinary Bladder</u>			
Hyperplasia	0	2(4%)	0
Dysplasia	0	1(2%)	0
Neoplasm NOS	0	0	1(2%)
Neoplasm NOS, Malignant	0	1(2%)	0
Carcinoma NOS	0	0	1(2%)
Squamous-Cell Carcinoma	0	21(45%)	43(90%)
Transitional-Cell Carcinoma	0	20(43%)	0
Sarcoma NOS	0	0	1(2%)
Leiomyosarcoma	0	2(4%)	2(4%)

The bladders of those few mice that did not have bladder neoplasms showed epithelial hyperplasia such as that seen in rats or dysplasia which was characterized by nuclear hyperchromatism and irregularity. In contrast to those in the rats, the bladder tumors in the mice more frequently invaded adjacent structures or metastasized to distant organs and were more frequently squamous in type rather than transitional-cell. The mouse bladder tumors tended to be large, often filling the lumen of the organs and obstructing the ureters. In consequence, kidneys of the tumor-bearing animals had some degree of hydronephrosis. As in the rats, transitional-cell tumors occurred in the renal pelvis of two p-cresidine-dosed females.

There were no tumors of the nasal cavity in control mice. Of the mice fed p-cresidine two developed sarcomas, one an adenocarcinoma, and one a squamous-cell carcinoma of the nasal cavity. However, no olfactory neuroblastomas were observed.

The incidence of hepatocellular carcinomas in low dose female mice (13/45 [29 percent]), but not in the high dose female group (6/43 [14 percent]), was increased relative to controls (0/46). The incidence of liver tumors in dosed male mice was not increased. The early mortality resulting from fulminating tumors of the bladder in both sexes may have obscured a dose-related response of the liver.

Based upon this pathologic examination, p-cresidine was carcinogenic in B6C3F1 mice because it induced carcinomas of the urinary bladder in both sexes and hepatocellular carcinomas in female mice.

#### D. Statistical Analyses of Results

The results of the statistical analyses of tumor incidence in mice are summarized in Tables 5 and 6. The analysis is included for every type of malignant tumor in either sex where at least two such tumors were observed in at least one of the control or p-cresidine-dosed groups and where such tumors were observed in at least 5 percent of the group. Because of the high mortality noted in mice of both sexes, these analyses were based on those mice surviving at least 52 weeks or at least until the development of the first tumor at the site of interest, whichever occurred first.

In both male and female mice the combined incidence of carcinomas NOS, squamous-cell carcinomas, or transitional-cell carcinomas of the urinary bladder was increased in the dosed groups compared to the control groups. These tumors were observed as early as week 44



TABLE 5

TIME-ADJUSTED ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT  
SPECIFIC SITES IN MALE MICE TREATED WITH p-CRESIDINE<sup>a,f</sup>

TOPOGRAPHY: MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Lung: Alveolar/Bronchiolar Carcinoma, or Alveolar/Bronchiolar Adenoma <sup>b</sup>	7/50(0.14)	2/39(0.05)	1/21(0.05)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	0.366	0.340
Lower Limit	---	0.039	0.008
Upper Limit	---	1.788	2.374
Weeks to First Observed Tumor	97	106	92
Hematopoietic System: Leukemia or Malignant Lymphoma <sup>b</sup>	4/50(0.08)	0/42(0.00)	0/22(0.00)
P Values <sup>c</sup>	P = 0.047(N)	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	0.000	0.000
Lower Limit	---	0.000	0.000
Upper Limit	---	1.278	2.370
Weeks to First Observed Tumor	97	---	---
Liver: Hepatocellular Carcinoma <sup>b,f</sup>	10/50(0.20)	11/41(0.27)	3/32(0.09)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	1.341	0.469
Lower Limit	---	0.574	0.088
Upper Limit	---	3.146	1.648
Weeks to First Observed Tumor	60	65	44

TABLE 5 (Continued)

TOPOGRAPHY:MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Liver: Hepatocellular Carcinoma, or Hepatocellular Adenoma <sup>b,f</sup>	15/50(0.30)	11/41(0.27)	3/32(0.09)
P Values <sup>c</sup>	P = 0.028(N)	N.S.	P = 0.024(N)
Relative Risk (Control) <sup>d</sup>	---	0.894	0.313
Lower Limit	---	0.418	0.062
Upper Limit	---	1.836	0.991
Weeks to First Observed Tumor	60	65	44
Urinary Bladder: Squamous-Cell Carcinoma of Transitional-Cell Carcinoma <sup>b,f</sup>	0/50(0.00)	40/42(0.95)	31/31(1.00)
P Values <sup>c</sup>	P < 0.001	P < 0.001	P < 0.001
Departure from Linear Trend <sup>e</sup>	P < 0.001	---	---
Relative Risk (Control) <sup>d</sup>	---	Infinite	Infinite
Lower Limit	---	17.734	20.738
Upper Limit	---	Infinite	Infinite
Weeks to First Observed Tumor	---	51	44

TABLE 5 (Concluded)

aTreated groups received time-weighted average doses of 0.22 or 0.46 percent in feed.

bNumber of tumor-bearing animals/number of animals examined at site (proportion).

cThe probability level for the Cochran-Armitage test is given beneath the incidence of tumors in the control group when  $P < 0.05$ ; otherwise, not significant (N.S.) is indicated. The probability level for the Fisher exact test for the comparison of a treated group with the control group is given beneath the incidence of tumors in the treated group when  $P < 0.05$ ; otherwise, not significant (N.S.) is indicated. For both Cochran-Armitage and Fisher exact tests a negative designation (N) indicates a lower incidence in the treated group(s) than in the control group.

dThe 95 percent confidence interval on the relative risk of the treated group to the control group.

eThe probability level of the test for departure from linear trend is given beneath the control group when  $P < 0.05$ .

fThese analyses were based solely upon animals surviving at least 52 weeks, except for sites where the first tumor of interest was observed earlier than 52 weeks in any group of this sex and species, where the analyses were based upon all animals that survived until or past the date that the first tumor was observed.

TIME-ADJUSTED ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT  
SPECIFIC SITES IN FEMALE MICE TREATED WITH P-CRESIDINE<sup>a, f</sup>

TOPOGRAPHY:MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Lung: Alveolar/Bronchiolar Carcinoma <sup>b</sup>	3/45(0.07)	3/45(0.07)	0.34(0.00)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	1.000	0.000
Lower Limit	---	0.141	0.000
Upper Limit	---	7.095	2.173
Weeks to First Observed Tumor	97	106	---
Lung: Alveolar/Bronchiolar Carcinoma or Alveolar/Bronchiolar Adenoma <sup>b</sup>	4/45(0.09)	4/45(0.09)	1/34(0.03)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	1.000	0.331
Lower Limit	---	0.198	0.007
Upper Limit	---	5.051	3.135
Weeks to First Observed Tumor	97	106	70
Hematopoietic System: Leukemia or Malignant Lymphoma <sup>b</sup>	6/46(0.13)	10/45(0.22)	1/36(0.03)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Departure from Linear Trend <sup>e</sup>	P = 0.024	---	---
Relative Risk (Control) <sup>d</sup>	---	1.704	0.213
Lower Limit	---	0.615	0.005
Upper Limit	---	5.229	1.638
Weeks to First Observed Tumor	69	87	54

TABLE 6 (Continued)

TOPOGRAPHY: MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Liver: Hepatocellular Carcinoma <sup>b,f</sup>	0/45(0.00)	13/44(0.30)	6/40(0.15)
P Values <sup>c</sup>	P = 0.028	P < 0.001	P = 0.009
Departure from Linear Trend <sup>e</sup>	P < 0.001	---	---
Relative Risk (Control) <sup>d</sup>	---	Infinitive	Infinitive
Lower Limit	---	4.126	1.809
Upper Limit	---	Infinitive	Infinitive
Weeks to First Observed Tumor	---	53	44
Liver: Hepatocellular Carcinoma, or Hepatocellular Adenoma <sup>b,f</sup>	0/45(0.00)	14/44(0.32)	6/40(0.15)
P Values <sup>c</sup>	P = 0.030	P < 0.001	P = 0.009
Departure from Linear Trend <sup>e</sup>	P < 0.001	---	---
Relative Risk (Control) <sup>d</sup>	---	Infinitive	Infinitive
Lower Limit	---	4.484	1.809
Upper Limit	---	Infinitive	Infinitive
Weeks to First Observed Tumor	---	53	44
Urinary Bladder: Carcinoma NOS, Squamous-Cell Carcinoma, or Transitional-Cell Carcinoma <sup>b,f</sup>	0/45(0.00)	41/46(0.89)	44/46(0.96)
P Values <sup>c</sup>	P < 0.001	P < 0.001	P < 0.001
Departure from Linear Trend <sup>e</sup>	P < 0.001	---	---
Relative Risk (Control) <sup>d</sup>	---	Infinitive	Infinitive
Lower Limit	---	14.258	16.126
Upper Limit	---	Infinitive	Infinitive
Weeks to First Observed Tumor	---	53	40



TABLE 6 (Continued)

TOPOGRAPHY:MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Urinary Bladder: Transitional-Cell Carcinoma, Neoplasm NOS, Neoplasm NOS Malignant, Carcinoma NOS, or Squamous-Cell Carcinoma <sup>b,f</sup>	0/45(0.00)	42/46(0.91)	45/46(0.98)
P Values <sup>c</sup>	P < 0.001	P < 0.001	P < 0.001
Departure from Linear Trend <sup>e</sup>	P < 0.001	---	---
Relative Risk (Control) <sup>d</sup>	---	Infinit	Infinit
Lower Limit	---	14.784	17.153
Upper Limit	---	Infinit	Infinit
Weeks to First Observed Tumor	---	53	40
Thyroid: Follicular-Cell Adenoma <sup>b</sup>	1/28(0.04)	2/31(0.06)	0/30(0.00)
P Values <sup>c</sup>	N.S.	N.S.	N.S.
Relative Risk (Control) <sup>d</sup>	---	1.806	0.000
Lower Limit	---	0.100	0.000
Upper Limit	---	103.026	17.204
Weeks to First Observed Tumor	97	94	---

TABLE 6 (Concluded)

aTreated groups received time-weighted average doses of 0.22 or 0.44 percent in feed.

bNumber of tumor-bearing animals/number of animals examined at site (proportion).

cThe probability level for the Cochran-Armitage test is given beneath the incidence of tumors in the control group when  $P < 0.05$ ; otherwise, not significant (N.S.) is indicated. The probability level for the Fisher exact test for the comparison of a treated group with the control group is given beneath the incidence of tumors in the treated group when  $P < 0.05$ ; otherwise, not significant (N.S.) is indicated. For both Cochran-Armitage and Fisher exact tests a negative designation (N) indicates a lower incidence in the treated group(s) than in the control group.

dThe 95 percent confidence interval on the relative risk of the treated group to the control group.

eThe probability level of the test for departure from linear trend is given beneath the control group when  $P < 0.05$ .

fThese analyses were based solely upon animals surviving at least 52 weeks, except for sites where the first tumor of interest was observed earlier than 52 weeks in any group of this sex and species, where the analyses were based upon all animals that survived until or past the date that the first tumor was observed.

in the males and week 40 in the females. For both sexes, the Cochran-Armitage test indicated a significant ( $P < 0.001$ ) positive association between dose and incidence. These were supported by significant ( $P < 0.001$ ) Fisher exact test results for both dose levels compared to the control.

In female mice an increased incidence of hepatocellular carcinomas was observed in the dosed groups compared to the control. The Cochran-Armitage test indicated a significant ( $P = 0.028$ ) positive association between dose and incidence. This was supported by significant Fisher exact test results of both the high ( $P = 0.009$ ) and low ( $P < 0.001$ ) dose groups to the control. The departure from linear trend was also significant ( $P = 0.001$ ) since tumor incidence was greater in the low dose group than in the high dose group. In historical control data compiled by this laboratory for the NCI Carcinogenesis Testing Program 19/275 (7 percent) of the female untreated control B6C3F1 mice had a hepatocellular carcinoma or a hepatocellular adenoma. For the male mice, however, the Cochran-Armitage test indicated a significant ( $P = 0.028$ ) negative association between dose and the combined incidence of hepatocellular carcinomas and hepatocellular adenomas. The Fisher exact test comparing high dose to control also indicated a significant ( $P = 0.024$ ) negative association. The historical control rate for untreated male B6C3F1 mice was 88/275 (32 percent) with these lesions. It was considered,

however, that the shorter survival in the high dose than in the control and low dose groups accounted for the apparently negative trend.

Based on these results the administration of p-cresidine was associated with the increased incidence of hepatocellular carcinoma in female mice and of carcinomas of the urinary bladder in both male and female mice.

## V. DISCUSSION

Mortality rates were dose-related for both sexes of both species. Accelerated mortality of high dose animals during the latter half of the study was associated with bladder tumors. That incidences of certain tumors were higher in low dose than in high dose groups was probably due to this accelerated mortality.

Addition of p-cresidine to the diet was associated with increased incidences of neoplasms at several sites in rats. Male rats, having a higher incidence of neoplasms than females in the urinary bladder, nasal cavity, liver, and brain, appeared to be more sensitive to the tumorigenic effects of p-cresidine.

The urinary bladder was a target organ of p-cresidine carcinogenicity in rats. Bladder neoplasms were observed in dosed rats, but not in their controls. For both male and female rats the incidence of carcinomas (combined incidence of papillary carcinomas, squamous-cell carcinomas, transitional-cell papillomas, transitional-cell carcinomas and undifferentiated carcinomas) of the urinary bladder was dose-related and was significantly higher in each dosed group than in the corresponding control group. An elevated incidence of hyperplasia was also observed in urinary bladders of low dose rats of both sexes.

Dose-related increases in the incidence of neoplasms were also observed in nasal cavities of rats. The incidence of olfactory neuroblastomas in females and the combined incidence of neuroblastoma



and olfactory neuroblastoma of the nasal cavity in males were both significantly associated with p-cresidine dosage. The incidence of these neoplasms in high dose groups of both sexes was significantly higher than in controls. The combined incidences of carcinomas and adenocarcinomas of the nasal cavity or nasopharynx were significantly dose-related in male rats.

Among low dose male rats, the incidence of liver tumors (the combined incidence of neoplastic nodules, hepatocellular carcinomas, or mixed hepato/cholangio carcinomas) was significantly higher than in controls.

The urinary bladder was a target organ for carcinogenicity in mice as well as in rats. Although the spectrum of dose-related bladder neoplasms in mice was similar to that in rats, the bladder tumors in mice were more invasive and more frequently metastasized. In both male and female mice, the incidence of carcinomas (combined incidence of carcinomas NOS, squamous-cell carcinomas, and transitional-cell carcinomas) of the urinary bladder was significantly associated with p-cresidine dosage and was significantly greater in each dosed group than in controls. The bladders of those few mice without bladder neoplasms showed evidence of hyperplasia or dysplasia. No bladder neoplasms were found in control mice.

The incidence of hepatocellular carcinomas in each dosed female mouse group was significantly greater than in controls, and this

increased incidence was significantly associated with concentration of p-cresidine in the diet.

Under the conditions of this bioassay, p-cresidine was carcinogenic to Fischer 344 rats, causing increased incidences of carcinomas and of papillomas of the urinary bladder in both sexes, increased incidences of olfactory neuroblastomas in both sexes, and of liver tumors (hepatocellular carcinomas, hepato/cholangio carcinomas and neoplastic nodules) in males. p-Cresidine was also found to be carcinogenic in B6C3F1 mice, causing carcinomas of the urinary bladders in both sexes and hepatocellular carcinomas in females.

## VI. BIBLIOGRAPHY

- Anthony, H.M., and G.M. Thomas, "Tumors of the Urinary Bladder: An Analysis of the Occupations of 1,030 Patients in Leeds, England." Journal of the National Cancer Institute 45:879-895, 1970.
- Armitage, P., Statistical Methods in Medical Research, Chapter 14. J. Wiley & Sons, New York, 1971.
- Berenblum, I., editor, Carcinogenicity Testing. International Union Against Cancer, Technical Report Series, Vol. 2. International Union Against Cancer, Geneva, 1969.
- Chemical Abstracts Service. The Chemical Abstracts Service (CAS) Ninth Collective Index, Volumes 76-85, 1972-1976. American Chemical Society, Washington, D.C., 1977.
- Compton, E.D., Group Director, Environmental Control, Chemicals, The Sherwin-Williams Company, Cleveland, Ohio. Letter to Dr. J. Donoso, The MITRE Corporation, METREK Division, McLean, Virginia, May 31, 1977.
- Cox, D.R., Analysis of Binary Data, Chapters 4 and 5. Methuen and Co., Ltd., London, 1970.
- Cox, D.R., "Regression Models and Life-Tables." Journal of the Royal Statistical Society, Series "B" 34:187-220, 1972.
- Gart, J.J., "The Comparison of Proportions: A Review of Significance Tests, Confidence Limits, and Adjustments for Stratification." International Statistical Institute Review 39:148-169, 1971.
- Kaplan, E.L., and P. Meier, "Nonparametric Estimation from Incomplete Observations." Journal of the American Statistical Association 53:457-481, 1958.
- Linhart, M.S., J.A. Cooper, R.L. Martin, N.P. Page, and J.A. Peters, "Carcinogenesis Bioassay Data System." Computers and Biomedical Research 7:230-248, 1974.
- Mehta, N., Sadtler Research Institute, Philadelphia, Pennsylvania. Personal communication, 24 August 1978.
- Miller, R.G., Simultaneous Statistical Inference. McGraw-Hill Book Co., New York, 1966.

- Pollock, J.R.A. and R. Stevens, editors, Dictionary of Organic Compounds, 4th edition. Volume 1. Oxford University Press, New York, 1965.
- Saffiotti, U., R. Montesano, A.R. Sellakumar, F. Cefis, and D.G. Kaufman, "Respiratory Tract Carcinogenesis in Hamsters Induced by Different Numbers of Administration of Benzo (a) Pyrene and Ferric Oxide." Cancer Research 32:1073-1079, 1972.
- Society of Dyers and Colourists. Colour Index, 3rd edition, Volume 3. Yorkshire, England, p. 4706, 1971.
- Squire, R.A. and M.H. Levitt, "Report of a Workshop on Classification of Specific Hepatocellular Lesions in Rats." Cancer Research 35:3214-3223, 1975.
- Tarone, R.E., "Tests for Trend in Life-Table Analysis." Biometrika 62:679-682, 1975.
- Urso, S., Research Analyst, Chemical-Environmental Program, Chemical Industries Center, Stanford Research Institute, Menlo Park, California. Personal communication, June 8, 1977.
- Wynder, E.L., J. Onderdonk, and N. Mantel, "An Epidemiological Investigation of Cancer of the Bladder." Cancer 16:1388-1407, 1963.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100



APPENDIX A

SUMMARY OF THE INCIDENCE OF NEOPLASMS  
IN RATS TREATED WITH p-CRESIDINE

10

TABLE A1  
SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE  
RATS TREATED WITH p-CRESIDINE

	CONTROL (UNTR) 01-0220	LOW DOSE 01-0260	HIGH DOSE 01-0265
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS NECROPSIED	48	50	47
ANIMALS EXAMINED HISTOPATHOLOGICALLY**	48	49	47
INTEGUMENTARY SYSTEM			
*SKIN	(48)	(50)	(47)
TRICHOEPITHELIOMA		2 (4%)	
*SUBCUT TISSUE	(48)	(50)	(47)
FIBROMA	3 (6%)	2 (4%)	1 (2%)
FIBROSARCOMA	2 (4%)	2 (4%)	
LIPOSARCOMA		1 (2%)	
RESPIRATORY SYSTEM			
*NASAL CAVITY	(48)	(50)	(47)
NEOPLASM, NOS, MALIGNANT			1 (2%)
CARCINOMA, NOS			1 (2%)
ADENOMA, NOS		1 (2%)	
NEUROBLASTOMA			1 (2%)
OLFACTORY NEUROBLASTOMA		1 (2%)	20 (43%)
*NASOPHARYNX	(48)	(50)	(47)
ADENOCARCINOMA, NOS			1 (2%)
*LUNG	(48)	(49)	(45)
HEPATOCELLULAR CARCINOMA, METAST			1 (2%)
ALVEOLAR/BRONCHIOLAR ADENOMA	3 (6%)		
FIBROSARCOMA, METASTATIC	1 (2%)		
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(48)	(50)	(47)
MALIGNANT LYMPHOMA, NOS	1 (2%)		1 (2%)
UNDIFFERENTIATED LEUKEMIA	2 (4%)	1 (2%)	
*SPLEEN	(48)	(49)	(45)
MYELOMONOCYTIC LEUKEMIA	4 (8%)		

\* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

\* NUMBER OF ANIMALS NECROPSIED

\*\* EXCLUDES PARTIALLY AUTOLYZED ANIMALS

TABLE A1  
(CONTINUED)

	CONTROL (UNTR) 01-0220	LOW DOSE 01-0260	HIGH DOSE 01-0265
#LYMPH NODE MALIGNANT LYMPHOMA, NOS	(43)	(48)	(38) 1 (3%)
#SUBMANDIBULAR L.NODE OLFACTORY NEUROBLASTOMA, METASTA	(43)	(48)	(38) 1 (3%)
CIRCULATORY SYSTEM			
NONE			
DIGESTIVE SYSTEM			
#SALIVARY GLAND ACINAR-CELL ADENOMA	(45)	(48) 1 (2%)	(43)
#PAROTID GLAND OLFACTORY NEUROBLASTOMA, METASTA	(45)	(48)	(43) 1 (2%)
#LIVER NEOPLASTIC NODULE HEPATOCELLULAR CARCINOMA MIXED HEPATO/CHOLANGIO CARCINOMA	(48)	(49) 10 (20%) 2 (4%) 1 (2%)	(46) 1 (2%) 1 (2%)
#STOMACH SQUAMOUS CELL PAPILLOMA BASAL-CELL CARCINOMA	(48)	(49) 2 (4%) 1 (2%)	(46)
URINARY SYSTEM			
#KIDNEY/PELVIS TRANSITIONAL-CELL PAPILLOMA TRANSITIONAL-CELL CARCINOMA, INV	(48)	(49) 1 (2%) 1 (2%)	(46)
#URINARY BLADDER UNDIFFERENTIATED CARCINOMA PAPILLARY CARCINOMA SQUAMOUS CELL CARCINOMA TRANSITIONAL-CELL PAPILLOMA TRANSITIONAL-CELL CARCINOMA LEIOMYOSARCOMA	(48)	(48)  1 (2%) 14 (29%) 15 (31%)	(47) 1 (2%) 4 (9%) 6 (13%) 3 (6%) 31 (66%) 2 (4%)
*PROSTATIC URETHRA UNDIFFERENTIATED CARCINOMA, INVA	(48)	(50)	(47) 1 (2%)

# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY  
\* NUMBER OF ANIMALS NECROPSIED

TABLE A1  
(CONTINUED)

	CONTROL (UNTR) 01-0220	LOW DOSE 01-0260	HIGH DOSE 01-0265
ENDOCRINE SYSTEM			
*PITUITARY	(45)	(48)	(38)
ADENOMA, NOS	10 (22%)	1 (2%)	3 (8%)
CHROMOPHOBE ADENOMA		4 (8%)	
BASOPHIL ADENOMA		4 (8%)	
*ADRENAL	(46)	(48)	(44)
CORTICAL ADENOMA		3 (6%)	
PHEOCHROMOCYTOMA	4 (9%)	7 (15%)	3 (7%)
*THYROID	(43)	(41)	(40)
FOLLICULAR-CELL ADENOMA		1 (2%)	
C-CELL ADENOMA	3 (7%)	1 (2%)	1 (3%)
C-CELL CARCINOMA	2 (5%)	1 (2%)	1 (3%)
*PARATHYROID	(25)	(21)	(15)
ADENOMA, NOS	1 (4%)		
*PANCREATIC ISLETS	(44)	(48)	(40)
ISLET-CELL ADENOMA		1 (2%)	2 (5%)
ISLET-CELL CARCINOMA		1 (2%)	
REPRODUCTIVE SYSTEM			
*PREPUTIAL GLAND	(48)	(50)	(47)
CARCINOMA, NOS		2 (4%)	1 (2%)
PAPILLOMA, NOS			1 (2%)
ADENOMA, NOS		1 (2%)	
*PROSTATE	(45)	(48)	(46)
TRANSITIONAL-CELL CARCINOMA, INV		1 (2%)	
*TESTIS	(48)	(47)	(46)
INTERSTITIAL-CELL TUMOR	37 (77%)	45 (96%)	23 (50%)
NERVOUS SYSTEM			
*BRAIN/MENINGES	(46)	(48)	(43)
HEPATOCELLULAR CARCINOMA, METAST			1 (2%)
*BRAIN	(46)	(48)	(43)
NEOPLASM, NOS, METASTATIC			1 (2%)
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			



TABLE A1  
(CONTINUED)

	CONTROL (UNTR) 01-0220	LOW DOSE 01-0260	HIGH DOSE 01-0265
CARCINOMA, NOS, METASTATIC			1 (2%)
OSTEOSARCOMA, METASTATIC	1 (2%)		
GLIOMA, NOS			2 (5%)
OLFACTORY NEUROBLASTOMA, INVASIV			2 (5%)
OLFACTORY NEUROBLASTOMA, METASTA			5 (12%)
#BRAIN STEM GLIOMA, NOS	(46)	(48)	(43) 1 (2%)
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
*SKULL	(48)	(50)	(47)
OSTEOSARCOMA	1 (2%)		
BOOY CAVITIES			
*BOOY CAVITIES	(48)	(50)	(47)
MESOTHELIOMA, NOS	1 (2%)	1 (2%)	2 (4%)
ALL OTHER SYSTEMS			
*MULTIPLE ORGANS	(48)	(50)	(47)
C-CELL CARCINOMA, METASTATIC		1 (2%)	
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	50	50	50
NATURAL DEATH	8	2	25
MORIBUND SACRIFICE	5	9	21
SCHEDULED SACRIFICE	5		
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	32	39	4
ANIMAL MISSING			
2 INCLUDES AUTOLYZED ANIMALS			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A1  
(CONCLUDED)

	CONTROL (UNTR) 01-0220	LOW DOSE 01-0260	HIGH DOSE 01-0265
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	42	49	46
TOTAL PRIMARY TUMORS	74	131	116
TOTAL ANIMALS WITH BENIGN TUMORS	40	48	28
TOTAL BENIGN TUMORS	61	91	37
TOTAL ANIMALS WITH MALIGNANT TUMORS	11	21	46
TOTAL MALIGNANT TUMORS	12	29	76
TOTAL ANIMALS WITH SECONDARY TUMORS#	2	3	11
TOTAL SECONDARY TUMORS	2	3	14
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT	1	11	2
TOTAL UNCERTAIN TUMORS	1	11	3
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

TABLE A2  
SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE  
RATS TREATED WITH p-CRESIDINE

	CONTROL (UNTR) 02-0220	LOW DOSE 02-0260	HIGH DOSE 02-0265
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS NECROPSIED	50	50	49
ANIMALS EXAMINED HISTOPATHOLOGICALLY**	50	49	48
INTEGUMENTARY SYSTEM			
*SKIN	(50)	(50)	(49)
SEBACEOUS ADENOCARCINOMA		1 (2%)	
FIBROMA		1 (2%)	
*SUBCUT TISSUE	(50)	(50)	(49)
SARCOMA, NOS	1 (2%)	1 (2%)	
FIBROSARCOMA		1 (2%)	
RESPIRATORY SYSTEM			
*NASAL CAVITY	(50)	(50)	(49)
SQUAMOUS CELL CARCINOMA			1 (2%)
ADENOCARCINOMA, NOS			1 (2%)
OLFACTORY NEUROBLASTOMA			11 (22%)
*LUNG	(50)	(48)	(48)
SQUAMOUS CELL CARCINOMA, METASTA		1 (2%)	
TRANSITIONAL-CELL CARCINOMA, MET		1 (2%)	1 (2%)
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(50)	(50)	(49)
MALIGNANT LYMPHOMA, NOS	2 (4%)		
*SPLEEN	(50)	(49)	(47)
MYELOMONOCYTIC LEUKEMIA	2 (4%)		
CIRCULATORY SYSTEM			
NONE			

\* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

\* NUMBER OF ANIMALS NECROPSIED

\*\* EXCLUDES PARTIALLY AUTOLYZED ANIMALS

TABLE A2  
(CONTINUED)

	CONTROL (UNTR) 02-0220	LOW DOSE 02-0260	HIGH DOSE 02-0265
DIGESTIVE SYSTEM			
*MOUTH	(50)	(50)	(49)
SQUAMOUS CELL PAPILLOMA			1 (2%)
#SALIVARY GLAND	(46)	(48)	(43)
ACINAR-CELL ADENOMA		1 (2%)	
#LIVER	(50)	(48)	(48)
NEOPLASTIC NODULE		4 (8%)	
#PANCREAS	(46)	(49)	(43)
TRANSITIONAL-CELL CARCINOMA, MET			1 (2%)
#STOMACH	(49)	(48)	(46)
SQUAMOUS CELL CARCINOMA			1 (2%)
URINARY SYSTEM			
#KIDNEY	(49)	(49)	(46)
UNDIFFERENTIATED CARCINOMA			1 (2%)
#KIDNEY/PELVIS	(49)	(49)	(46)
TRANSITIONAL-CELL CARCINOMA		1 (2%)	
#URINARY BLADDER	(47)	(49)	(46)
NEOPLASM, NOS			2 (4%)
PAPILLARY CARCINOMA			1 (2%)
SQUAMOUS CELL CARCINOMA		9 (18%)	7 (15%)
TRANSITIONAL-CELL PAPILLOMA		6 (12%)	2 (4%)
TRANSITIONAL-CELL CARCINOMA		22 (45%)	33 (72%)
LEIOMYOSARCOMA		2 (4%)	3 (7%)
*URETHRA	(50)	(50)	(49)
SQUAMOUS CELL PAPILLOMA		1 (2%)	
ENDOCRINE SYSTEM			
#PITUITARY	(40)	(45)	(40)
ADENOMA, NOS	16 (40%)	2 (4%)	6 (15%)
CHROMOPHOBE ADENOMA	1 (3%)	9 (20%)	1 (3%)
#ADRENAL	(48)	(48)	(44)
CORTICAL ADENOMA		5 (10%)	1 (2%)

# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY  
\* NUMBER OF ANIMALS NECROPSIED

TABLE A2  
(CONTINUED)

	CONTROL (UNTR) 02-0220	LOW DOSE 02-0260	HIGH DOSE 02-0265
PHEOCHROMOCYTOMA	6 (13%)	1 (2%)	1 (2%)
*THYROID	(43)	(42)	(40)
FOLLICULAR-CELL CARCINOMA		2 (5%)	
C-CELL ADENOMA		4 (10%)	
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND	(50)	(50)	(49)
ADENOCARCINOMA, NOS	1 (2%)		
FIBROADENOMA	6 (12%)	4 (8%)	1 (2%)
*CLITORAL GLAND	(50)	(50)	(49)
CARCINOMA, NOS		2 (4%)	
ADENOMA, NOS		1 (2%)	
*UTERUS	(48)	(47)	(45)
ADENOCARCINOMA, NOS	1 (2%)		
FIBROMA	2 (4%)		
ENDOMETRIAL STROMAL POLYP	2 (4%)	8 (17%)	4 (9%)
ENDOMETRIAL STROMAL SARCOMA	2 (4%)		
CARCINOSARCOMA		1 (2%)	
*UTERUS/ENDOMETRIUM	(48)	(47)	(45)
ADENOCARCINOMA, NOS		1 (2%)	
ADENOC/SQUAMOUS METAPLASIA		1 (2%)	
NERVOUS SYSTEM			
*BRAIN	(50)	(48)	(46)
SQUAMOUS CELL CARCINOMA, METASTA			2 (4%)
ADENOCARCINOMA, NOS, INVASIVE			1 (2%)
GLIOMA, NOS		1 (2%)	
OLIGODENDROGLIOMA		1 (2%)	
OLFACTORY NEUROBLASTOMA, INVASIV			1 (2%)
OLFACTORY NEUROBLASTOMA, METASTA			3 (7%)
*CEREBRAL CORTEX	(50)	(48)	(46)
OLFACTORY NEUROBLASTOMA, METASTA			1 (2%)
SPECIAL SENSE ORGANS			
*EAR CANAL	(50)	(50)	(49)
SQUAMOUS CELL CARCINOMA			1 (2%)

\* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY  
\* NUMBER OF ANIMALS NECROPSIED



TABLE A2  
(CONTINUED)

	CONTROL(UNTR) 02-0220	LOW DOSE 02-0260	HIGH DOSE 02-0265
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
*ABDOMINAL CAVITY	(50)	(50)	(49)
TRANSITIONAL-CELL CARCINOMA, INV			1 (2%)
ALL OTHER SYSTEMS			
NONE			
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	50	50	50
NATURAL DEATH‡	7	3	30
MORIBUND SACRIFICE	2	6	20
SCHEDULED SACRIFICE	5		
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	36	41	
ANIMAL MISSING			
‡ INCLUDES AUTOLYZED ANIMALS			
‡ NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A2  
(CONCLUDED)

	CONTROL (UNTR) 02-0220	LOW DOSE 02-0260	HIGH DOSE 02-0265
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	27	48	44
TOTAL PRIMARY TUMORS	42	93	79
TOTAL ANIMALS WITH BENIGN TUMORS	23	35	17
TOTAL BENIGN TUMORS	33	43	17
TOTAL ANIMALS WITH MALIGNANT TUMORS	9	36	42
TOTAL MALIGNANT TUMORS	9	46	60
TOTAL ANIMALS WITH SECONDARY TUMORS*		2	9
TOTAL SECONDARY TUMORS		2	11
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT		4	2
TOTAL UNCERTAIN TUMORS		4	2
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
* SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

APPENDIX B

SUMMARY OF THE INCIDENCE OF NEOPLASMS  
IN MICE TREATED WITH p-CRESIDINE

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

TABLE B1  
SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE  
MICE TREATED WITH p-CRESIDINE

	CONTROL (UNTR) 05-0220	LOW DOSE 05-0260	HIGH DOSE 05-0265
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS MISSING		1	3
ANIMALS NECROPSIED	50	47	41
ANIMALS EXAMINED HISTOPATHOLOGICALLY **	50	43	37
INTEGUMENTARY SYSTEM			
NONE			
RESPIRATORY SYSTEM			
*NASAL CAVITY	(50)	(47)	(41)
ADENOCARCINOMA, NOS		1 (2%)	
SARCOMA, NOS			1 (2%)
FIBROSARCOMA		1 (2%)	
#LUNG	(50)	(40)	(37)
SQUAMOUS CELL CARCINOMA, METASTA		4 (10%)	2 (5%)
TRANSITIONAL-CELL CARCINOMA, MET		1 (3%)	
HEPATOCELLULAR CARCINOMA, METAST		2 (5%)	
ALVEOLAR/BRONCHIOLAR ADENOMA	5 (10%)	1 (3%)	
ALVEOLAR/BRONCHIOLAR CARCINOMA	2 (4%)	1 (3%)	1 (3%)
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(50)	(47)	(41)
MALIG.LYMPHOMA, HISTIOCYTIC TYPE	1 (2%)		
#SPLEEN	(50)	(37)	(33)
HEMANGIOMA	1 (2%)		
HEMANGIOSARCOMA	2 (4%)	1 (3%)	
*MESENTERIC L. NODE	(44)	(26)	(16)
MALIGNANT LYMPHOMA, NOS	1 (2%)		
MALIG.LYMPHOMA, HISTIOCYTIC TYPE	1 (2%)		
#LIVER	(50)	(41)	(37)
MALIG.LYMPHOMA, HISTIOCYTIC TYPE	1 (2%)		
CIRCULATORY SYSTEM			
NONE			
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			
** EXCLUDES PARTIALLY AUTOLYZED ANIMALS			



TABLE B1  
(CONTINUED)

	CONTROL (UNTR) 05-0220	LOW DOSE 05-0260	HIGH DOSE 05-0265
DIGESTIVE SYSTEM			
*LIVER	(50)	(41)	(37)
HEPATOCELLULAR ADENOMA	5 (10%)		
HEPATOCELLULAR CARCINOMA	10 (20%)	11 (27%)	3 (8%)
*PANCREAS	(50)	(30)	(31)
SQUAMOUS CELL CARCINOMA, INVASIV		1 (3%)	
*DUODENAL GLAND	(50)	(30)	(29)
ADENOMA, NOS		1 (3%)	
URINARY SYSTEM			
*URINARY BLADDER	(50)	(42)	(35)
SQUAMOUS CELL CARCINOMA		27 (64%)	28 (80%)
TRANSITIONAL-CELL CARCINOMA		13 (31%)	3 (9%)
LEIOMYOSARCOMA		2 (5%)	4 (11%)
*PROSTATIC URETHRA	(50)	(47)	(41)
SQUAMOUS CELL CARCINOMA, INVASIV		2 (4%)	
TRANSITIONAL-CELL CARCINOMA, INV		1 (2%)	
ENDOCRINE SYSTEM			
*THYROID	(39)	(31)	(29)
FOLLICULAR-CELL ADENOMA	1 (3%)		
FOLLICULAR-CELL CARCINOMA	1 (3%)		
REPRODUCTIVE SYSTEM			
*PROSTATE	(41)	(37)	(26)
SQUAMOUS CELL CARCINOMA, INVASIV		6 (16%)	2 (8%)
TRANSITIONAL-CELL CARCINOMA, INV		3 (8%)	1 (4%)
LEIOMYOSARCOMA, INVASIVE		1 (3%)	1 (4%)
*SEMINAL VESICLE	(50)	(47)	(41)
TRANSITIONAL-CELL CARCINOMA, INV		1 (2%)	
*COAGULATING GLAND	(50)	(47)	(41)
SQUAMOUS CELL CARCINOMA, INVASIV		1 (2%)	

\* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY  
\* NUMBER OF ANIMALS NECROPSIED

TABLE B1  
(CONTINUED)

	CONTROL (UNTR) 05-0220	LOW DOSE 05-0260	HIGH DOSE 05-0265
TRANSITIONAL-CELL CARCINOMA, INV		1 (2%)	
*TESTIS	(50)	(41)	(37)
EMBRYONAL CARCINOMA	1 (2%)		
NERVOUS SYSTEM			
*BRAIN	(50)	(36)	(32)
SARCOMA, NOS, METASTATIC			1 (3%)
SPECIAL SENSE ORGANS			
*HARDERIAN GLAND	(50)	(47)	(41)
CYSTADENOMA, NOS	1 (2%)	1 (2%)	
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
*ABDOMINAL CAVITY	(50)	(47)	(41)
SQUAMOUS CELL CARCINOMA, INVASIV		4 (9%)	3 (7%)
SQUAMOUS CELL CARCINOMA, METASTA		1 (2%)	
TRANSITIONAL-CELL CARCINOMA, INV		1 (2%)	
LEIOMYOSARCOMA, INVASIVE			1 (2%)
*PERITONEAL CAVITY	(50)	(47)	(41)
SQUAMOUS CELL CARCINOMA, INVASIV			2 (5%)
ALL OTHER SYSTEMS			
*MULTIPLE ORGANS	(50)	(47)	(41)
SQUAMOUS CELL CARCINOMA, INVASIV		4 (9%)	1 (2%)
SQUAMOUS CELL CARCINOMA, METASTA		2 (4%)	2 (5%)
TRANSITIONAL-CELL CARCINOMA, INV			2 (5%)
ADENOCARCINOMA, NOS, INVASIVE		1 (2%)	
LEIOMYOSARCOMA, INVASIVE		1 (2%)	

\* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY  
\* NUMBER OF ANIMALS NECROPSIED

TABLE BI  
(CONCLUDED)

	CONTROL (UNTR) 05-0220	LOW DOSE 05-0260	HIGH DOSE 05-0265
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	50	50	50
NATURAL DEATH <sup>a</sup>	3	37	40
MORIBUND SACRIFICE		5	7
SCHEDULED SACRIFICE	5		
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	42	7	
ANIMAL MISSING		1	3
<sup>a</sup> INCLUDES AUTOLYZED ANIMALS			
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	22	42	32
TOTAL PRIMARY TUMORS	33	60	40
TOTAL ANIMALS WITH BENIGN TUMORS	12	3	
TOTAL BENIGN TUMORS	13	3	
TOTAL ANIMALS WITH MALIGNANT TUMORS	16	42	32
TOTAL MALIGNANT TUMORS	20	57	40
TOTAL ANIMALS WITH SECONDARY TUMORS <sup>b</sup>		27	14
TOTAL SECONDARY TUMORS		38	18
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT			
TOTAL UNCERTAIN TUMORS			
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
<sup>b</sup> SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

TABLE B2  
SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE  
MICE TREATED WITH p-CRESIDINE

	CONTROL (UNTR) 06-0220	LOW DOSE 06-0260	HIGH DOSE 06-0265
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS MISSING	2		
ANIMALS NECROPSIED	47	47	48
ANIMALS EXAMINED HISTOPATHOLOGICALLY**	47	47	48
-----			
INTEGUMENTARY SYSTEM			
*SKIN	(47)	(47)	(48)
KERATOACANTHOMA	1 (2%)		
*SUBCUT TISSUE	(47)	(47)	(48)
FIBROSARCOMA		1 (2%)	
-----			
RESPIRATORY SYSTEM			
*NASAL CAVITY	(47)	(47)	(48)
SQUAMOUS CELL CARCINOMA			1 (2%)
*LUNG	(46)	(47)	(44)
SQUAMOUS CELL CARCINOMA, METASTA		1 (2%)	3 (7%)
HEPATOCELLULAR CARCINOMA, METAST		1 (2%)	
ALVEOLAR/BRONCHIOLAR ADENOMA	1 (2%)	1 (2%)	1 (2%)
ALVEOLAR/BRONCHIOLAR CARCINOMA	3 (7%)	3 (6%)	
-----			
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(47)	(47)	(48)
MALIGNANT LYMPHOMA, NOS	1 (2%)	1 (2%)	
MALIG. LYMPHOMA, UNDIFFER-TYPE		1 (2%)	
MALIG. LYMPHOMA, HISTIOCYTIC TYPE	3 (6%)		
MALIGNANT LYMPHOMA, MIXED TYPE		5 (11%)	
*SPLEEN	(45)	(44)	(38)
HEMANGIOSARCOMA		1 (2%)	
MALIG. LYMPHOMA, HISTIOCYTIC TYPE	2 (4%)		
MALIGNANT LYMPHOMA, MIXED TYPE		1 (2%)	
*MANDIBULAR L. NODE	(38)	(36)	(28)
HEMANGIOMA		1 (3%)	

\* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

\* NUMBER OF ANIMALS NECROPSIED

\*\* EXCLUDES PARTIALLY AUTOLYZED ANIMALS

TABLE B2  
(CONTINUED)

	CONTROL (UNTR) 06-0220	LOW DOSE 06-0260	HIGH DOSE 06-0265
*ABDOMINAL LYMPH NODE SQUAMOUS CELL CARCINOMA, METASTA	(38)	(38) 1 (3%)	(28)
*MESENTERIC L. NODE MALIG. LYMPHOMA, HISTIOCYTIC TYPE	(38)	(38) 1 (3%)	(28)
*LIVER MALIG. LYMPHOMA, HISTIOCYTIC TYPE	(46)	(45)	(43) 1 (2%)
MALIGNANT LYMPHOMA, MIXED TYPE		1 (2%)	
CIRCULATORY SYSTEM			
NONE			
DIGESTIVE SYSTEM			
*LIVER SQUAMOUS CELL CARCINOMA, METASTA	(46)	(45)	(43) 1 (2%)
HEPATOCELLULAR ADENOMA		1 (2%)	
HEPATOCELLULAR CARCINOMA		13 (29%)	6 (14%)
*STOMACH ADENOCARCINOMA, NOS	(46)	(40) 1 (3%)	(36)
URINARY SYSTEM			
*KIDNEY/PELVIS TRANSITIONAL-CELL CARCINOMA	(46)	(45) 1 (2%)	(45) 1 (2%)
*URINARY BLADDER NEOPLASM, NOS	(45)	(47)	(48) 1 (2%)
NEOPLASM, NOS, MALIGNANT		1 (2%)	
CARCINOMA, NOS			1 (2%)
SQUAMOUS CELL CARCINOMA		21 (45%)	43 (90%)
TRANSITIONAL-CELL CARCINOMA		20 (43%)	
SARCOMA, NOS			1 (2%)
LEIOMYOSARCOMA		2 (4%)	2 (4%)
ENDOCRINE SYSTEM			
*PITUITARY ADENOMA, NOS	(33) 1 (3%)	(32)	(26)

\* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY  
\* NUMBER OF ANIMALS NECROPSIED



TABLE B2  
(CONTINUED)

	CONTROL (UNTR) 06-0220	LOW DOSE 06-0260	HIGH DOSE 06-0265
*ADRENAL/CAPSULE ADENOMA, NOS	(40) 1 (3%)	(42)	(39)
*THYROID FOLLICULAR-CELL ADENOMA	(29) 1 (3%)	(32) 2 (6%)	(37)
REPRODUCTIVE SYSTEM			
*UTERUS HEMANGIOSARCOMA	(45)	(46) 1 (2%)	(36)
*OVARY PAPILLARY ADENOMA HEMANGIOMA	(40) 1 (3%)	(41) 1 (2%)	(35)
NERVOUS SYSTEM			
*BRAIN SQUAMOUS CELL CARCINOMA, INVASIV	(44)	(45)	(39) 1 (3%)
SPECIAL SENSE ORGANS			
*HARDERIAN GLAND PAPILLARY ADENOMA PAPILLARY ADENOCARCINOMA PAPILLARY CYSTADENOMA, NOS	(47)	(47) 1 (2%) 1 (2%)	(48) 1 (2%)
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
*ABDOMINAL CAVITY SQUAMOUS CELL CARCINOMA, INVASIV	(47)	(47) 2 (4%)	(48) 2 (4%)
*PERITONEAL CAVITY SQUAMOUS CELL CARCINOMA, INVASIV	(47)	(47)	(48) 2 (4%)
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE B2  
(CONCLUDED)

	CONTROL (UNTR) 06-0220	LOW DOSE 06-0260	HIGH DOSE 06-0265
ALL OTHER SYSTEMS			
*MULTIPLE ORGANS	(47)	(47)	(48)
SQUAMOUS CELL CARCINOMA, INVASIV			2 (4%)
SQUAMOUS CELL CARCINOMA, METASTA		1 (2%)	
LEIOMYOSARCOMA, INVASIVE		1 (2%)	
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	50	50	50
NATURAL DEATH <sup>a</sup>	4	21	41
MORIBUND SACRIFICE	3	4	8
SCHEDULED SACRIFICE	5		
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	36	25	1
ANIMAL MISSING	2		
<sup>a</sup> INCLUDES AUTOLYZED ANIMALS			
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	14	45	45
TOTAL PRIMARY TUMORS	15	83	59
TOTAL ANIMALS WITH BENIGN TUMORS	5	7	2
TOTAL BENIGN TUMORS	6	7	2
TOTAL ANIMALS WITH MALIGNANT TUMORS	9	45	44
TOTAL MALIGNANT TUMORS	9	76	56
TOTAL ANIMALS WITH SECONDARY TUMORS <sup>#</sup>		6	9
TOTAL SECONDARY TUMORS		7	11
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT			1
TOTAL UNCERTAIN TUMORS			1
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
<sup>#</sup> SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

## APPENDIX C

SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC  
LESIONS IN RATS TREATED WITH p-CRESIDINE



TABLE C1  
SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS  
IN MALE RATS TREATED WITH p-CRESIDINE

	CONTROL (UNTR) 01-0220	LOW DOSE 01-0260	HIGH DOSE 01-0265
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS NECROPSIED	48	50	47
ANIMALS EXAMINED HISTOPATHOLOGICALLY **	48	49	47
INTEGUMENTARY SYSTEM			
*SKIN	(48)	(50)	(47)
EPIDERMAL INCLUSION CYST	1 (2%)		
*SUBCUT TISSUE	(48)	(50)	(47)
ABSCCESS, NOS	3 (6%)		
RESPIRATORY SYSTEM			
*NASAL CAVITY	(48)	(50)	(47)
HYPERPLASIA, EPITHELIAL			5 (11%)
POLYP		1 (2%)	
*NASAL TURBINATE	(48)	(50)	(47)
INFLAMMATION, SUPPURATIVE	1 (2%)		
#LUNG/BRONCHUS	(48)	(49)	(45)
CALCIFICATION, NOS			1 (2%)
#LUNG	(48)	(49)	(45)
BRONCHOPNEUMONIA, ACUTE			1 (2%)
INFLAMMATION, ACUTE FOCAL			2 (4%)
PNEUMONIA, CHRONIC MURINE		3 (6%)	
METAPLASIA, NOS			2 (4%)
#LUNG/ALVEOLI	(48)	(49)	(45)
CALCIFICATION, FOCAL			1 (2%)
HEMATOPOIETIC SYSTEM			
#SPLEEN	(48)	(49)	(45)
INFARCT, NOS	1 (2%)		
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY * NUMBER OF ANIMALS NECROPSIED ** EXCLUDES PARTIALLY AUTOLYZED ANIMALS			



TABLE C1  
(CONTINUED)

	CONTROL (UNTR) 01-0220	LOW DOSE 01-0260	HIGH DOSE 01-0265
ATROPHY, NOS HEMATOPOIESIS		1 (2%)	1 (2%)
*THYMUS HYPERPLASIA, NOS	(32) 1 (3%)	(33)	(23)
CIRCULATORY SYSTEM			
*HEART PERIARTERITIS CALCIFICATION, FOCAL	(48)	(49) 1 (2%)	(45) 1 (2%)
*MYOCARDIUM DEGENERATION, NOS CALCIFICATION, FOCAL	(48)	(49) 6 (12%)	(45) 1 (2%) 2 (4%)
*BLOOD VESSEL MEDIAL CALCIFICATION	(48)	(50)	(47) 2 (4%)
*CORONARY ARTERY CALCIFICATION, FOCAL	(48)	(50)	(47) 1 (2%)
DIGESTIVE SYSTEM			
*SALIVARY GLAND HYPERPLASIA, INTRAUCTAL DYSPLASIA, NOS	(45)	(48) 1 (2%)	(43) 1 (2%)
*LIVER METAMORPHOSIS FATTY BASOPHILIC CYTO CHANGE CLEAR-CELL CHANGE HYPERPLASIA, FOCAL	(48) 2 (4%)  3 (6%)	(49) 5 (10%) 1 (2%) 1 (2%)	(46) 1 (2%) 1 (2%) 1 (2%)
*PANCREAS INFLAMMATION, NOS	(44) 2 (5%)	(48)	(40)
*PANCREATIC ACINUS ATROPHY, NOS	(44)	(48)	(40) 1 (3%)
*STOMACH ULCER, NOS CALCIFICATION, FOCAL	(48)	(49) 1 (2%)	(46) 1 (2%) 2 (4%)

\* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY  
\* NUMBER OF ANIMALS NECROPSIED

TABLE C1  
(CONTINUED)

	CONTROL (UNTR) 01-0220	LOW OOSE 01-0260	HIGH OOSE 01-0265
HYPERPLASIA, BASAL CELL		8 (16%)	
*GASTRIC MUCOSA CALCIFICATION, NOS	(48)	(49)	(46) 1 (2%)
*GASTRIC MUSCULARIS INFLAMMATION, NECROTIZING CALCIFICATION, NOS	(48)	(49)	(46) 1 (2%) 1 (2%)
URINARY SYSTEM			
*KIDNEY	(48)	(49)	(46)
HYORONEPHROSIS		1 (2%)	3 (7%)
HEMORRHAGE			1 (2%)
GLOMERULONEPHRITIS, NOS			3 (7%)
INFLAMMATION, INTERSTITIAL			1 (2%)
NEPHROPATHY	35 (73%)		
NEPHROSIS, NOS		26 (53%)	14 (30%)
NECROSIS, MEUOLLARY			3 (7%)
CALCIFICATION, NOS			3 (7%)
CALCIFICATION, FOCAL			3 (7%)
HYPERPLASIA, TUBULAR CELL			1 (2%)
HYPERPLASIA, EPITHELIAL			2 (4%)
*KIDNEY/MEOUILLA AMYLOIOOSIS	(48)	(49)	(46) 1 (2%)
*RENAL PAPILLA INFLAMMATION, ACUTE NECROTIZING	(48)	(49)	(46) 1 (2%)
*KIDNEY/TUBULE CAST, NOS	(48)	(49)	(46) 1 (2%)
FIBROSIS			1 (2%)
DEGENERATION, NOS			1 (2%)
CALCIFICATION, FOCAL			1 (2%)
*KIDNEY/PELVIS HYPERPLASIA, EPITHELIAL	(48)	(49) 1 (2%)	(46)
*URINARY BLADOER INFLAMMATION, ACUTE FOCAL	(48)	(48)	(47) 1 (2%)
HYPERPLASIA, EPITHELIAL		11 (23%)	1 (2%)
HYPERPLASIA, PAPILLARY			1 (2%)
POLYP, INFLAMMATORY			3 (6%)

\* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY  
\* NUMBER OF ANIMALS NECROPSIED

TABLE CI  
(CONTINUED)

	CONTRDL (UNTR) 01-0220	LDW DDSE 01-0260	HIGH DOSE 01-0265
ENDOCRINE SYSTEM			
#PITUITARY	(45)	(48)	(38)
HYPERPLASIA, NODULAR		1 (2%)	
HYPERPLASIA, FOCAL	1 (2%)		
HYPERPLASIA, BASOPHILIC		1 (2%)	1 (3%)
#PITUITARY/BASOPHIL	(45)	(48)	(38)
HYPERPLASIA, NODULAR		1 (2%)	
#ADRENAL CORTEX	(46)	(48)	(44)
HYPERPLASIA, FOCAL		1 (2%)	
#ADRENAL MEDULLA	(46)	(48)	(44)
HYPERPLASIA, NOS	1 (2%)		
HYPERPLASIA, FOCAL	3 (7%)		
#PANCREATIC ISLETS	(44)	(48)	(40)
HYPERPLASIA, NOS	1 (2%)		
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND	(48)	(50)	(47)
HEMORRHAGIC CYST		1 (2%)	
#PROSTATE	(45)	(48)	(46)
INFLAMMATIDN, ACUTE			5 (11%)
INFLAMMATION, ACUTE FOCAL			1 (2%)
INFLAMMATIDN, ACUTE/CHRONIC			2 (4%)
*SEMINAL VESICLE	(48)	(50)	(47)
ATROPHY, NOS			1 (2%)
#TESTIS	(48)	(47)	(46)
MINERALIZATION	1 (2%)		
ATROPHY, NOS	4 (8%)		2 (4%)
HYPERPLASIA, INTERSTITIAL CELL	3 (6%)		2 (4%)
#TESTIS/TUBULE	(48)	(47)	(46)
DEGENERATION, NOS			7 (15%)
*EPIIDYMISS	(48)	(50)	(47)
ABSCESS, NOS	1 (2%)		

# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

\* NUMBER OF ANIMALS NECROPSIED

TABLE C1  
(CONCLUDED)

	CONTROL (UNTR) 01-0220	LOW DOSE 01-0260	HIGH DOSE 01-0265
NERVOUS SYSTEM			
*BRAIN GLIOSIS	(46)	(48) 1 (2%)	(43)
SPECIAL SENSE ORGANS			
*EYE CATARACT	(48) 1 (2%)	(50)	(47)
*EYE/RETINA DEGENERATION, NOS ATROPHY, NOS	(48) 2 (4%)	(50) 1 (2%)	(47)
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
*MULTIPLE ORGANS MINERALIZATION	(48)	(50) 1 (2%)	(47)
SPECIAL MORPHOLOGY SUMMARY			
AUTO/NECROPSY/HISTO PERF	1		
AUTO/NECROPSY/NO HISTO		1	
AUTOLYSIS/NO NECROPSY	2		3
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE C2  
SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS  
IN FEMALE RATS TREATED WITH p-CRESIDINE

	CONTROL(UNTR) 02-0220	LOW DOSE 02-0260	HIGH DOSE 02-0265
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS NECROPSIED	50	50	49
ANIMALS EXAMINED HISTOPATHOLOGICALLY**	50	49	48
-----			
INTEGUMENTARY SYSTEM			
NONE			
-----			
RESPIRATORY SYSTEM			
*NASAL CAVITY	(50)	(50)	(49)
HYPERPLASIA, EPITHELIAL			8 (16%)
*NASAL TURBINATE	(50)	(50)	(49)
INFLAMMATION, NOS	1 (2%)		
*LUNG	(50)	(48)	(48)
INFLAMMATION, NOS	1 (2%)		
INFLAMMATION, INTERSTITIAL	1 (2%)		
ABSCCESS, NOS	2 (4%)		
PNEUMONIA, CHRONIC MURINE	1 (2%)		
GRANULOMA, NOS	1 (2%)		
-----			
HEMATOPOIETIC SYSTEM			
*BONE MARROW	(48)	(46)	(43)
OSTEOSCLEROSIS			2 (5%)
*SPLEEN	(50)	(49)	(47)
HEMATOPOIESIS	7 (14%)		
-----			
CIRCULATORY SYSTEM			
*HEART	(50)	(48)	(47)
ENDOCARDIOSIS		1 (2%)	
*MYOCARDIUM	(50)	(48)	(47)
FIBROSIS, FOCAL			1 (2%)

\* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

\* NUMBER OF ANIMALS NECROPSIED

\*\* EXCLUDES PARTIALLY AUTOLYZED ANIMALS

TABLE C2  
(CONTINUED)

	CONTROL (UNTR) 02-0220	LOW DOSE 02-0260	HIGH DOSE 02-0265
DIGESTIVE SYSTEM			
*SALIVARY GLAND DYSPLASIA, NOS	(46)	(48) 1 (2%)	(43)
*LIVER	(50)	(48)	(48)
INFLAMMATION, FOCAL		1 (2%)	
METAMORPHOSIS FATTY	4 (8%)		2 (4%)
BASOPHILIC CYTO CHANGE			1 (2%)
HYPERPLASIA, FOCAL	9 (18%)	1 (2%)	1 (2%)
*LIVER/CENTRIOLOBULAR NECROSIS, NOS	(50)	(48) 1 (2%)	(48)
*LIVER/HEPATOCYTES HYPERPLASIA, FOCAL	(50)	(48)	(48) 2 (4%)
*STOMACH	(49)	(48)	(46)
ULCER, NOS			1 (2%)
HYPERPLASIA, PAPILLARY	1 (2%)		
HYPERPLASIA, BASAL CELL		2 (4%)	
*GASTRIC MUCOSA ULCER, NOS	(49)	(48)	(46) 1 (2%)
*DUODENUM ULCER, NOS	(47)	(49) 1 (2%)	(46)
URINARY SYSTEM			
*KIDNEY	(49)	(49)	(46)
HYDRONEPHROSIS			2 (4%)
GLOMERULONEPHRITIS, NOS	1 (2%)		2 (4%)
PYELONEPHRITIS, FOCAL			1 (2%)
PYELONEPHRITIS SUPPURATIVE			1 (2%)
PYELONEPHRITIS, CHRONIC		1 (2%)	
NEPHROPATHY	18 (37%)		
NEPHROSIS, NOS		2 (4%)	1 (2%)
NECROSIS, MEDULLARY			2 (4%)
HYPERPLASIA, EPITHELIAL			3 (7%)
*KIDNEY/MEDULLA CALCIFICATION, FOCAL	(49)	(49)	(46) 1 (2%)
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			



TABLE C2  
(CONTINUED)

	CONTROL (UNTR) 02-0220	LOW DOSE 02-0260	HIGH DOSE 02-0265
*RENAL PAPILLA CALCIFICATION, FOCAL	(49)	(49)	(46) 1 (2%)
*KIDNEY/TUBULE CALCIFICATION, FOCAL	(49)	(49)	(46) 1 (2%)
*KIDNEY/PELVIS INFLAMMATION, ACUTE/CHRONIC CALCIFICATION, FOCAL	(49)	(49)	(46) 1 (2%) 1 (2%)
*URINARY BLADDER INFLAMMATION, ACUTE INFLAMMATION, ACUTE FOCAL HYPERPLASIA, EPITHELIAL HYPERPLASIA, PAPILLARY POLYP, INFLAMMATORY	(47)	(49)  10 (20%) 1 (2%)	(46) 2 (4%) 1 (2%) 1 (2%) 5 (11%)
ENDOCRINE SYSTEM			
*PITUITARY MINERALIZATION HYPERPLASIA, NOS	(40) 1 (3%)	(45) 1 (2%)	(40)
*ZONA GLOMERULOSA METAMORPHOSIS FATTY	(48)	(48)	(44) 1 (2%)
*THYROID HYPERPLASIA, C-CELL	(43) 1 (2%)	(42)	(40)
*PARATHYROID HYPERPLASIA, NODULAR	(31) 1 (3%)	(28)	(22)
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND GALACTOCELE HYPERPLASIA, NOS	(50) 2 (4%) 1 (2%)	(50)	(49)
*UTERUS ABSCCESS, NOS NECROSIS, NOS	(48) 1 (2%) 1 (2%)	(47)	(45)
*UTERUS/ENDOMETRIUM INFLAMMATION, NOS	(48) 1 (2%)	(47)	(45)

\* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY  
\* NUMBER OF ANIMALS NECROPSIED

TABLE C2  
(CONCLUDED)

	CONTROL (UNTR) 02-0220	LOW DOSE 02-0260	HIGH DOSE 02-0265
INFLAMMATION, ACUTE			1 (2%)
*OVARY	(49)	(47)	(43)
INFLAMMATION, NOS	1 (2%)		
ABSCCESS, NOS		1 (2%)	
DEGENERATION, CYSTIC	2 (4%)		
NERVOUS SYSTEM			
*BRAIN	(50)	(48)	(46)
HYDROCEPHALUS, NOS			1 (2%)
GLIOSIS			1 (2%)
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
*ABDOMINAL CAVITY	(50)	(50)	(49)
NECROSIS, FAT		1 (2%)	
ALL OTHER SYSTEMS			
NONE			
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	11		
AUTO/NECROPSY/HISTO PERF	1		
AUTO/NECROPSY/NO HISTO		1	1
AUTOLYSIS/NO NECROPSY			1
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100

APPENDIX D

SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC  
LESIONS IN MICE TREATED WITH p-CRESIDINE



TABLE D1  
SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS  
IN MALE MICE TREATED WITH p-CRESIDINE

	CONTROL (UNTR) 0S-0220	LOW DOSE 0S-0260	HIGH DOSE 0S-026S
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS MISSING		1	3
ANIMALS NECROPSIED	50	47	41
ANIMALS EXAMINED HISTOPATHOLOGICALLY**	50	43	37
INTEGUMENTARY SYSTEM			
*SUBCUT TISSUE ABSCCESS, NOS	(50) 1 (2%)	(47)	(41)
RESPIRATORY SYSTEM			
*LUNG/BRONCHUS INFLAMMATION, NOS	(50) 1 (2%)	(40)	(37)
HEMATOPOIETIC SYSTEM			
*SPLEEN ATROPHY, NOS	(50)	(37) 17 (46%)	(33)
*MESENTERIC L. NODE HEMATOPOIESIS	(44) 1 (2%)	(26)	(16)
CIRCULATORY SYSTEM			
*MYOCARDIUM MINERALIZATION DEGENERATION, NOS	(50)	(40) 1 (3%) 1 (3%)	(36)
*CORONARY ARTERY MEDIAL CALCIFICATION	(50)	(47)	(41) 1 (2%)
*PULMONARY ARTERY MEDIAL CALCIFICATION	(50)	(47)	(41) 1 (2%)
DIGESTIVE SYSTEM			
*LIVER NECROSIS, COAGULATIVE	(50) 1 (2%)	(41)	(37)

\* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

\* NUMBER OF ANIMALS NECROPSIED

\*\* EXCLUDES PARTIALLY AUTOLYZED ANIMALS



TABLE D1  
(CONTINUED)

	CONTROL (UNTR) 05-0220	LOW DOSE 05-0260	HIGH DOSE 05-0265
*PANCREATIC ACINUS ATROPHY, NOS HYPERTROPHY, FOCAL	(50) 1 (2%) 1 (2%)	(30)	(31)
*GASTRIC MUCOSA CALCIFICATION, NOS	(50)	(32)	(33) 1 (3%)
*GASTRIC MUSCULARIS CALCIFICATION, NOS	(50)	(32)	(33) 1 (3%)
URINARY SYSTEM			
*KIDNEY HYDRONEPHROSIS	(50) 1 (2%)	(43) 29 (67%)	(36) 31 (86%)
*URINARY BLADDER HYPERPLASIA, EPITHELIAL DYSPLASIA, NOS	(50)	(42)	(35) 3 (9%) 1 (3%)
ENDOCRINE SYSTEM			
*PANCREATIC ISLETS INFLAMMATION, NOS HYPERPLASIA, ADENOMATOUS	(50) 3 (6%) 1 (2%)	(30)	(31)
REPRODUCTIVE SYSTEM			
*TESTIS ATROPHY, NOS	(50) 1 (2%)	(41)	(37)
*TESTIS/TUBULE MINERALIZATION	(50) 2 (4%)	(41)	(37)
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
NONE			
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE D1  
(CONCLUDED)

	CONTROL (UNTR) 05-0220	LOW DOSE 05-0260	HIGH DOSE 05-0265
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
*PERITONEUM INFLAMMATION, ACUTE/CHRONIC	(50)	(47)	(41) 1 (2%)
ALL OTHER SYSTEMS			
*MULTIPLE ORGANS ATROPHY, NOS	(50)	(47) 1 (2%)	(41)
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	22	1	1
ANIMAL MISSING/NO NECROPSY		1	3
NECROPSY PERF/NO HISTO PERFORMED			2
AUTO/NECROPSY/NO HISTO		4	2
AUTOLYSIS/NO NECROPSY		2	6
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE D2  
SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS  
IN FEMALE MICE TREATED WITH p-CRESIDINE

	CONTROL (UNTR) 06-0220	LOW DOSE 06-0260	HIGH DOSE 06-0265
ANIMALS INITIALLY IN STUDY	50	50	50
ANIMALS MISSING	2		
ANIMALS NECROPSIED	47	47	48
ANIMALS EXAMINED HISTOPATHOLOGICALLY **	47	47	48
-----			
INTEGUMENTARY SYSTEM			
*SUBCUT TISSUE INFLAMMATION, NOS	(47) 1 (2%)	(47)	(48)
-----			
RESPIRATORY SYSTEM			
#LUNG	(46)	(47)	(44)
THROMBOSIS, NOS		1 (2%)	
INFLAMMATION, ACUTE NECROTIZING		1 (2%)	
-----			
HEMATOPOIETIC SYSTEM			
*BONE MARROW	(39)	(44)	(39)
MYELOFIBROSIS		14 (32%)	
MYELOPOIESIS			3 (8%)
*SPLEEN	(45)	(44)	(38)
ATROPHY, NOS		9 (20%)	2 (5%)
HYPERPLASIA, RETICULUM CELL		2 (5%)	
HYPERPLASIA, LYMPHOID		2 (5%)	
HEMATOPOIESIS	1 (2%)		2 (5%)
*MEDIASTINAL L.N.OOE	(38)	(38)	(28)
HYPERPLASIA, LYMPHOID		1 (3%)	
-----			
CIRCULATORY SYSTEM			
*HEART	(45)	(47)	(42)
PERIARTERITIS			1 (2%)
*MYOCARDIUM	(45)	(47)	(42)
CALCIFICATION, NOS			1 (2%)

# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

\* NUMBER OF ANIMALS NECROPSIED

\*\* EXCLUDES PARTIALLY AUTOLYZED ANIMALS

TABLE D2  
(CONTINUED)

	CONTROL (UNTR) 06-0220	LOW DOSE 06-0260	HIGH DOSE 06-0265
*ARTERY INFLAMMATION, NOS	(47)	(47) 1 (2%)	(48)
*CORONARY ARTERY MEDIAL CALCIFICATION	(47)	(47)	(48) 1 (2%)
*MEDIASTINAL ARTERY MEDIAL CALCIFICATION	(47)	(47)	(48) 1 (2%)
DIGESTIVE SYSTEM			
*LIVER	(46)	(45)	(43)
ABSCCESS, CHRONIC			1 (2%)
METAMORPHOSIS FATTY	1 (2%)	2 (4%)	
ANGIECTASIS		1 (2%)	
*PANCREAS	(45)	(42)	(33)
CYSTIC DUCTS		1 (2%)	
*PANCREATIC ACINUS	(45)	(42)	(33)
ATROPHY, NOS		1 (2%)	
*STOMACH	(46)	(40)	(36)
ULCER, NOS			1 (3%)
HYPERKERATOSIS		1 (3%)	
*GASTRIC MUCOSA	(46)	(40)	(36)
INFLAMMATION, ACUTE FOCAL		1 (3%)	
CALCIFICATION, FOCAL		1 (3%)	
*GASTRIC MUSCULARIS	(46)	(40)	(36)
CALCIFICATION, FOCAL			1 (3%)
URINARY SYSTEM			
*KIDNEY	(46)	(45)	(45)
HYDRONEPHROSIS		18 (40%)	37 (82%)
GLOMERULONEPHRITIS, NOS	1 (2%)	1 (2%)	
PYELONEPHRITIS, ACUTE		1 (2%)	1 (2%)
PYELONEPHRITIS, CHRONIC			1 (2%)
NEPHROPATHY		1 (2%)	
NEPHROSIS, NOS			1 (2%)

# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY  
\* NUMBER OF ANIMALS NECROPSIED

TABLE D2  
(CONTINUED)

	CONTROL (UNTR) 06-0220	LOW DOSE 06-0260	HIGH DOSE 06-0265
GLOMERULOSCLEROSIS, NOS INFARCT, NOS CALCIFICATION, FOCAL		1 (2%)	1 (2%) 1 (2%)
*URINARY BLADDER HYPERPLASIA, EPITHELIAL DYSPLASIA, NOS	(45)	(47) 2 (4%) 1 (2%)	(48)
ENDOCRINE SYSTEM			
NONE			
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND HYPERPLASIA, NOS	(47) 1 (2%)	(47)	(48)
*UTERUS HYDROMETRA	(45)	(46) 1 (2%)	(36)
*UTERUS/ENDOMETRIUM INFLAMMATION, ACUTE HYPERPLASIA, CYSTIC	(45)	(46) 1 (2%) 14 (30%)	(36) 1 (3%)
*OVARY/OVIDUCT DEGENERATION, NOS	(45) 1 (2%)	(46)	(36)
*OVARY THROMBOSIS, NOS PLASMA-CELL INFILTRATE	(40)	(41) 1 (2%) 1 (2%)	(35)
NERVOUS SYSTEM			
*BRAIN HYDROCEPHALUS, NOS	(44)	(45) 1 (2%)	(39)
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
NONE			
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY * NUMBER OF ANIMALS NECROPSIED			

TABLE D2  
(CONCLUDED)

	CONTROL (UNTR) 06-0220	LOW DOSE 06-0260	HIGH DOSE 06-0265
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
*MULTIPLE ORGANS PERIARTERITIS	(47)	(47)	(48) 1 (2%)
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	30		1
ANIMAL MISSING/NO NECROPSY	2		
AUTO/NECROPSY/HISTO PERF	1		1
AUTOLYSIS/NO NECROPSY	1	3	2
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			





Review of the Bioassay of *p*-Cresidine\* for Carcinogenicity  
by the Data Evaluation/Risk Assessment Subgroup  
of the Clearinghouse on Environmental Carcinogens

June 29, 1978

The Clearinghouse on Environmental Carcinogens was established in May, 1976, in compliance with DHEW Committee Regulations and the Provisions of the Federal Advisory Committee Act. The purpose of the Clearinghouse is to advise the Director of the National Cancer Institute (NCI) on its bioassay program to identify and to evaluate chemical carcinogens in the environment to which humans may be exposed. The members of the Clearinghouse have been drawn from academia, industry, organized labor, public interest groups, State health officials, and quasi-public health and research organizations. Members have been selected on the basis of their experience in carcinogenesis or related fields and, collectively, provide expertise in chemistry, biochemistry, biostatistics, toxicology, pathology, and epidemiology. Representatives of various Governmental agencies participate as ad hoc members. The Data Evaluation/Risk Assessment Subgroup of the Clearinghouse is charged with the responsibility of providing a peer review of reports prepared on NCI-sponsored bioassays of chemicals studied for carcinogenicity. It is in this context that the below critique is given on the bioassay of *p*-Cresidine for carcinogenicity.

The reviewer agreed with the conclusion in the report that *p*-Cresidine was carcinogenic in both rats and mice, under the conditions of test. He said that the nature of the tumors observed clearly indicate that they were treatment-related. He noted that the *p*-Cresidine was dissolved in acetone but that there was no acetone control group. This shortcoming, however, did not interfere with the conclusion on the carcinogenicity of *p*-Cresidine. Based on the results of the study, he said that *p*-Cresidine would appear to pose a carcinogenic risk to humans. The reviewer moved that the report on the bioassay of *p*-Cresidine be accepted as written. The motion was approved without objection.

Clearinghouse Members present:

Arnold L. Brown (Chairman), Mayo Clinic

Paul Nettesheim, National Institute of Environmental  
Health Sciences

Verne Ray, Pfizer Medical Research Laboratory

Verald K. Rowe, Dow Chemical U.S.A.

Michael B. Shimkin, University of California at San Diego

Louise Strong, University of Texas Health Sciences Center

\* Subsequent to this review, changes may have been made in the bioassay report either as a result of the review or other reasons. Thus, certain comments and criticisms reflected in the review may no longer be appropriate.

ARTICLE

The first part of the paper deals with the general principles of the subject, and the second part with the details of the method. The first part is divided into two sections, the first of which deals with the general principles of the subject, and the second with the details of the method. The second part is divided into two sections, the first of which deals with the general principles of the subject, and the second with the details of the method.

The first part of the paper deals with the general principles of the subject, and the second part with the details of the method. The first part is divided into two sections, the first of which deals with the general principles of the subject, and the second with the details of the method. The second part is divided into two sections, the first of which deals with the general principles of the subject, and the second with the details of the method.

The first part of the paper deals with the general principles of the subject, and the second part with the details of the method. The first part is divided into two sections, the first of which deals with the general principles of the subject, and the second with the details of the method. The second part is divided into two sections, the first of which deals with the general principles of the subject, and the second with the details of the method.

Library  
National Institute of Health  
Bethesda, Maryland 20014

## DATE DUE

[illegible]

**NIH** LIBRARY  
Amazing Research

Amazing Resea. .  
Amazing Help.

<http://nihlibrary.nih.gov>

**10 Center Drive  
Bethesda, MD 20892-1150  
301-496-1080**

**GAYLORD**

PRINTED IN U.S.A.

NIH LIBRARY



4 0118 6487

